



The Impact of Orthodontic Forces on the Occurrence of Iatrogenic Tooth Root Resorption

Nino Tsilosani, Professor
Ana Galogre, PhD, DMD
Natia Natsvlishvili, Ph.D student
Grigol robakidze University
School of medicine, Georgia

[Doi: 10.19044/esipreprint.9.2022.p479](https://doi.org/10.19044/esipreprint.9.2022.p479)

Approved: 21 September 2022
Posted: 23 September 2022

Copyright 2022 Author(s)
Under Creative Commons BY-NC-ND
4.0 OPEN ACCESS

Cite As:

Tsilosani N., Galogre A. & Natsvlishvili N. (2022). *The Impact of Orthodontic Forces on the Occurrence of Iatrogenic Tooth Root Resorption*. ESI Preprints.
<https://doi.org/10.19044/esipreprint.9.2022.p479>

Abstract

Modern orthodontic treatment ensures optimal functional and aesthetic results, but does not exclude the risk of various complications, including external or internal resorption of tooth roots. One of the causes of root resorption is the impact of orthodontic forces during treatment with both non-removable (brace-system) and removable (aligners) appliances. The aforementioned complication of the treatment of occlusal anomalies, which occurs with a fairly high frequency, potentially threatens not only the patient's health, but also complicates the treatment outcomes and worsens the prognosis. The lack of pathognomonic symptoms of root resorption increases the difficulty of an accurate diagnosis. According to the literature, the issue of root resorption is very relevant in clinical dentistry but has not been studied in detail, and contradictory opinions can be found.

Keywords: Orthodontic forces, tooth root resorption, non-removable appliances (brace-system), removable devices (aligners)

Introduction

Root resorption (RR) in deciduous teeth is a normal physiologic response, resulting in exfoliation of the deciduous teeth with replacement by permanent dentition. However, the process of root resorption in permanent dentition has a pathologic basis. However, in permanent dentition, root resorption has a pathologic basis, and the etiology requires two phases: an injury and stimulus. Types of root resorption in permanent teeth may be broadly categorized into internal and external resorption. The RR process may be self-limiting and go undetected clinically; however, once initiated and with the initial injury and stimulus sustained, destruction of hard dental tissue will continue, and tooth tissue loss may occur. External RR has various causes and is more prevalent than internal RR, which is relatively rare. Several classifications have been proposed, with distinctive terms and categories used to describe tooth root resorption (Henrique et al., 2018).

Modern orthodontic treatment ensures optimal functional and aesthetic results, but does not exclude the risk of various complications, including external or internal resorption of tooth roots (Yuan et al.,(2020), Jianru et al., 2018, João et al., (2018), Caroline et al., (2015), Ahu et al., (2012)). Caroline The aforementioned complication of the treatment of occlusal anomalies, which occurs with a fairly high frequency, potentially threatens not only the patient's health, but also complicates the treatment outcomes and worsens the prognosis (Caroline et al., (2015)). The trauma received before orthodontic treatment, bone density and morphotype, the shape of tooth roots, patient's age, duration of treatment, mechanism of orthodontic treatment, equal distribution of forces on teeth, metabolic disorders, and others are considered to be the determining factors of root resorption.



Fig.1. Treatment with Brace system (Dr. Natia Natsvlshvili)



Fig.2. Teeth straightening with Clear Aligners (Dr. Natia Natsvlshvili)




Fig 3. Active extreme external apical resorption in maxillary and mandibular incisors, associated with orthodontic treatment (Alberto Consolaro, <https://doi.org/10.1590/2177-6709.24.5.020-028.oin>)

Classification of RR

Several classifications have been proposed, with distinctive terms and categories used to describe RR. Amongst all classifications, the Andreasen classification is fundamental and the most widespread, as all subsequent classifications have cited it, and comparisons amongst classifications are always made using this one. Moreover, the authors consider the Kanas classification to be the most complete with full descriptions and categorizations (Henrique et al., 2018). (Table.4).

Table.4. Classification of root resorption

Andreasen J.	1970	Localization, type, and trauma	<p>I. Internal resorption-Replacement, Inflammatory</p> <p>II. External resorption-Surface, Replacement, Inflammatory</p>
Kanas and Kanas	2011	Localization, aetiology and Type	<p>1. Dental Origin -Internal resorption-Infective, inflammatory, trauma: a) Radial pulp enlargement resorption; b) Metaplastic (replacement) pulp resorption</p> <p>2. Dental Original- External resorption-</p> <p>a) Physiological apical resorption: External surface resorption, TAR; b) Infective/Inflammatory resorption: Apical (pulp) inflammatory resorption, Cervical (periodontal) inflammatory resorption; c) Trauma (avulsed/luxated/fractured): TAB, Periapical replacement resorption (PARR) with ankylosis, Infective/inflammatory (apical or cervical); d) Pressure: Orthodontic (TAB, PARR without ankylosis), Impacted teeth, Occlusal forces; e) Idiopathic: Localized apical resorption (PARR without ankylosis), Multiple apical resorption (PARR without ankylosis); Multiple cervical resorption; f) Surgical: Bone grafts of alveolar clefts.</p> <p>Nondental Origin</p> <p>1. Internal resorption - Herpes zoster infection.</p> <p>2. External resorption - Neoplasia/cysts of the jaws, Systemic disorders.</p>

Diagnosis

The relevance of the problem was first voiced by Ottolengui R in 1914 (Jacob et al.,(2020)), since then many studies have been devoted to determining the cause-and-effect relationship between orthodontic treatment and root resorption, and the main research methods include radiological studies, the most informative of which is cone-beam computed tomography (CBCT) (Neeta et al, (2017), Jian-Hong et al.,(2013) that can enhance the diagnosis of resorptive lesions in earlier stages. The detailed visualization afforded by CBCT is also associated with an improvement in final outcomes.

The lack of pathognomonic symptoms of root resorption increases the difficulty of an accurate diagnosis. **An early diagnosis** is a crucial factor in the management of this pathology because the earlier the treatment is initiated, the less severe the resorption consequences will be. Sensibility tests might be useful for RR recognition (Jian-Hong et al.,(2013), Topkara et al.,(2012)) and are mandatory for differential diagnosis. Full extra- and intraoral clinical examination must be implemented as a first step.

Literature review

Today, the genetic and molecular aspects of this problem are actively researched. However, despite this, the issue has not been studied in detail, and contradictory opinions can be found (Krishnan et al.,(2017)). Although most studies do not confirm the correlation between the patient's age and the incidence of root resorption, according to Sameshima and Sinclair, the majority of cases occur in older patients and there is a statistically insignificant increase in the incidence of resorption in men. Geraldo and co-authors associate the mentioned tendency in men with the morphological peculiarities of the root and explain this by the pipette-like shape of the roots of the central incisors. Kathryn and Begole studied the roots of patients from four ethnic groups (Asians, Caucasians, Spanish, and African Americans) prior to orthodontic treatment. According to these authors, women in all four ethnic groups had significantly shorter roots compared to men, and when it came to ethnic differences, Caucasians and Spanish had longer roots compared to Asians and African Americans. According to their own data, Asians are less prone to root resorption compared to Caucasians and Spanish. According to Kathryn and Begole, short roots (Asians, African Americans, and females) are more likely to be resorbed by orthodontic forces than roots with long or atypical roots (Sameshima et al., (2001)). According to the literature, root resorption is the second most common complication of orthodontic treatment after enamel white spots. The majority of authors believe that resorption is most often observed on the roots of upper incisors, although it is not clear which incisor

root is resorbed more often - central or lateral ((Krishnan et al.,(2017), Jairo (2013), Al-Qawasmi RA et al., (2004)).

In the frequency of detection of this pathology on the upper jaw, the incisors are followed by the molars, and then the incisors, as for the lower jaw, resorption is manifested more in the incisors, followed by the central and lateral incisors.

According to other data, we find opposite results - resorption on the lower jaw is most often seen in the central and lateral incisors, while in the molars of the lower jaw, resorption of the distal root is more marked. In accordance with the same authors, the molars of the lower jaw are followed by the molars of the upper jaw and the premolars of the lower jaw, and finally, the lowest rate of resorption is found in the premolars of the upper jaw (Krishnan et al., 2017).

A number of authors focus on the root apex shape abnormally and its relationship with resorption (Jairo et al (2013), Sherrard et al (2017)). In roots with different apex shapes (single apex, blunted, bifurcated, pointed, immature) the lowest frequency of resorption was observed in roots with blunted apex. Compared to roots with a single apex, resorption is more marked in the roots with an apical bifurcation, and much less in immature teeth. According to the data of Levander E, Bajka R and Malmgren O, blunt roots are more inclined to resorption (Al-Qawasmi RA et al ., (2004)). A positive correlation was found between root length and resorption—longer roots were more disposed to resorption than shorter ones (Mah Jet al ., (2004)). Data concerning the resorption of endodontically treated teeth are noteworthy (Jairo et al., (2013., Wickwire (2007), Pustulka et al., (2021)). According to the data of S. W. Spurrier et al., compared to vital teeth, dentin hardening after endodontic treatment provides increased resistance to root resorption, However, this hypothesis is contradicted by the studies of N.A. Wickwire (Wickwire, 2007), where no statistically significant difference was found in terms of resorption between vital and nonvital teeth, although it was noted that compared to intact teeth, resorption is more evident in endodontically treated teeth according to reports, root resorption is more common in adult orthodontic patients than in adolescents, and it is more common in women than in men, and resorption is characterized by a sluggish course in repeat orthodontic patients than in primary patients (Elif Dilara Seker et al ., (2021)).

The relationship between 1, 25-DHCC (Vit. D3), parathyroid hormone (PTH), Ca imbalance and root resorption has been determined, and endocrine disorders such as Hyperparathyroidism, Paget's disease, Hypophosphatemia, Hypothyroidism, Hypopituitarism, Hyperpituitarism has been considered as a cause of dental maldevelopment (Jacob et al., (2020)).

Thus, according to the literature, the issue of root resorption is very relevant in clinical dentistry. There is a lack of methodological approaches to studying the etiology, spread, diagnosis, treatment, and prevention of the mentioned pathology. Researchers mainly look at the issue of resorption from the individual aspects of endodontics, orthodontics, and periodontology.

Based on the aforementioned, we decided to study the occurrence of iatrogenic tooth root resorption in orthodontic practice thoroughly. Since June 2022 the research investigation is being conducted at the training-research and clinical center of Grigol Robakidze University “GRUNI DENTAL”. **The aim of the study** is to evaluate the prevalence and peculiarities of manifestation of root resorption caused by orthodontic treatment using non-removable (brace-system) and removable (aligner) devices for the treatment of malocclusion in children, adolescents, and adult patients.

The objectives of the planned research are the following:

- The study of the prevalence of internal and external root resorption of different groups of teeth in orthodontic patients at different stages of treatment (6- and 12 months after treatment) through cone-beam computed tomography;
- Development of systematization of root resorption caused by orthodontic appliances;
- Determination of the correlation between the manifestation of hard tissue resorption and the patient's age and gender, the shape of the apex and the length of the root, teeth vitality/devitality, and the duration of orthodontic treatment;
- Study of bone morphotype.

Exclusion Criteria are the following: Patients with systemic diseases or the use of any prescription drugs that might have an impact on the bone metabolism processes; patients with odontogenic acute or chronic apical periodontitis.

Materials and methods

The study will be conducted on 60 patients of different age groups:

1. Study groups: A - 12- to 17-years-old patients with non-removable orthodontic appliances, Subgroup B - 18- to 35-years-old patients with non-removable orthodontic appliances;
2. Control groups - A1 - 12- to 17-years-old patients with removable orthodontic appliances, subgroup B1 - 18- to 35-years-old patients with removable orthodontic appliances;

In order to achieve the goal of the study, patients of both the research and control groups will be subjected to orthopantomographic and cone-beam computed tomography studies; Registration of anamnesis data and orthopantomography will be used only in the pre-treatment stage, and 6 and 12 months after treatment will be used for research both radiovisiographic and cone-beam computerized tomographic research. Statistical processing and comparative analysis of the obtained results will be carried out. Systematization of root resorption caused by orthodontic devices will also be developed.

Conclusion

We consider the results of the current study should significantly simplify the treatment planning by taking into account the individual peculiarities of the patient and choosing an orthodontic appliance in order to prevent complications due to orthodontics, they will also make the results of treatment more predictable, and contribute its stable and safe outcome.

References:

1. Anna Jacob et al, 2020 , A Literature Review on Orthodontically Induced Root Resorption: The Aftermath of the Pursuit of an Attractive Smile, *European Journal of Molecular & Clinical Medicine*;
2. Ahu Topkara, Ali I Karaman, and Chung H Kau, 2012, Apical root resorption caused by orthodontic forces: A brief review and a long-term observation, *Eur J Dent.* Oct; 6(4): 445–453
3. Al-Qawasmi RA, Hartsfield JK, Everett ET, Weaver MR, et al. 2004; Root resorption associated with orthodontic force in IL-1Beta knockout mouse. *J Musculoskelet Neuronal Interact* 4: 383– 85
4. Aryal Neeta, Mao Jing, 2017, Root Resorption in Orthodontic Treatment: Scoping Review, *Orthodontic Journal of Nepal*, Vol. 7 No. 2, December
5. Caroline Pelagio Raick Maués¹, Rizomar Ramos do Nascimento , Oswaldo de Vasconcellos Vilella. 2015, Severe root resorption resulting from orthodontic treatment: Prevalence and risk factors, *Dental Press J Orthod.* Jan-Feb;20(1):52-8
6. Castro IO, Alencar AH, Valladares-Neto J, Estrela C. 2013, Apical root resorption due to orthodontic treatment detected by cone-beam computed tomography. *The Angle orthodontist.* ;83(2):196–203
7. Consolaro Alberto, <https://doi.org/10.1590/2177-6709.24.5.020-028.oin>
8. Durack C, Patel S, Davies J, Wilson R, Manocci F. 2011. Diagnostic accuracy of small volume cone-beam computed tomography and

- intraoral periapical radiography for the detection of simulated external inflammatory root resorption. *Int Endod J*; 44: 136– 47.
9. Elif Dilara Seker, Berza Sen Yilmaz, Ahmet Yagci, 2021, Evaluation of apical root resorption in Class III patients who received one- or two-phase orthodontic treatment APOS *Trends in Orthodontics* • Volume 11 • Issue 2 • April-June
 10. Jairo Curado de Freitas, Olavo César Porto Lyra, Ana Helena Gonçalves de Alencar, Carlos Estrela, 2013, Long-term evaluation of apical root resorption after orthodontic treatment using periapical radiography and cone-beam computed tomography, *Dental Press J. Orthod.* 18 (4) • Aug
 11. Jian-Hong, Yua, Kuang-2013, A cone-beam computed tomography study of orthodontic apical root resorption, *Journal of Dental Sciences*, Volume 8, Issue 1, March, Pages 74-79
 12. Henrique Aidos , Patrícia Diogo , João Miguel Santos , 2018 , Root Resorption Classifications: A Narrative Review and a Clinical Aid Proposal for Routine Assessment, *Eur Endod J*, Nov 21;3(3):134-145
 13. Jianru Yi, Jiani Xiao b, Yu Li, Xiaobing Li, Zhihe Zhao, 2018. External apical root resorption in non-extraction cases after clear aligner therapy or fixed orthodontic treatment. *Journal of Dental Sciences* 13, 48e53,
 14. João Dalto Viganó Pastro, Adriana Cândida Albuquerque Nogueira, 2018, Factors Associated to Apical Root Resorption after Orthodontic Treatment, *Open Dent J.*; 12: 331–339
 15. Katarzyna Pustułka, Agata Trzcionka, Arkadiusz Dziedzic, Dariusz Skiba, and Marta Tanasiewicz, 2021. The Radiological Assessment of Root Features and Periodontal Structures in Endodontically Treated Teeth Subjected to Forces Generated by Fixed Orthodontic Appliances. A Prospective, Clinical Cohort Study, *Clin. Med.*, 10
 16. Krishnan V, 2017, Root Resorption with Orthodontic Mechanics: Pertinent Areas Revisited, *Australian Dental Journal*; 62:(1 Suppl): 71–77
 17. Krishnan V. 2005, Critical issues concerning root resorption: A contemporary review: *World. J Orthod* ; 6: 30– 40.
 18. Mah J, Prasad N. 2004; Dentine phosphoproteins in gingival crevicular fluid during root resorption. *Eur J Orthod* 26: 25– 30
 19. Sherrard JF, Rossouw EP, Benson BW, Carrillo R, Buschang PH. 2010; Accuracy and reliability of tooth and root lengths measured on cone-beam computed tomographs. *Am J Orthod Dentofac Orthop* 137: S100– 108.

20. Sameshima GT, Sinclair PM , 2001) Predicting and preventing root resorption – Part II – Treatment factors. Am J Orthod Dentofac Orthop;119:511–515.
21. Ugnė Sadauskienė, Vilija Berlin, Orthodontic treatment with clear aligners and apical root Resorption, 2020, Vilnius University hospital Zalgiris clinic, Vilnius, Lithuania Journal of Medical Sciences. April 2, - Volume 8 | Issue
22. Yuan Li, Shiyong Deng, Li Mei, Zhengzheng Li, Xinyun Zhang, Chao Yang & Yu Li, 2020. Prevalence and severity of apical root resorption during orthodontic treatment with clear aligners and fixed appliances: a cone-beam computed tomography study, Progress in Orthodontics volume 21, Article number: 1;
23. Wickwire N.A., 2007, Orthodontic Root Resorption of Endodontically Treated Teeth, (February), Journal of Endodontics 33(2):119-22