

Petrographic and Geochemical Characteristics of the Djabatoure Massif Metamagmatites from Pan-African Orogen in Togo, West Africa

Sarakawa Abalo Malibida Kpanzou

Department of Geology, Faculty of Sciences, University of Lomé, Togo
Department of Civil Engineering, Laboratory of Geotechnics and Mining,
Regional Training Center for Road Maintenance
(CERFER - Entente Council), Lomé, Togo

Gnanwasou Alayi

Yao Agbossoumonde

Mahaman Sani Tairou

Department of Geology, Faculty of Sciences, University of Lomé, Togo

José María Gonzalez-Jimenez

Instituto Andaluz de Ciencias de la Tierra, CSIC-Universidad de Granada,
Avda. de las Palmeras 4, 18100 Armilla, Granada, Spain

Antonio Garcia-Casco

Departamento de Mineralogía y Petrología
Universidad de Granada, Granada, Spain

[Doi: 10.19044/esipreprint.7.2023.p564](https://doi.org/10.19044/esipreprint.7.2023.p564)

Approved: 27 July 2023

Posted: 29 July 2023

Copyright 2023 Author(s)

Under Creative Commons BY-NC-ND

4.0 OPEN ACCESS

Cite As:

Malibida Kpanzou S.A., Alayi G., Agbossoumonde Y. Tairou M.S., Gonzalez-Jimenez J.M. & Garcia-Casco A. (2023). *Petrographic and Geochemical Characteristics of the Djabatoure Massif Metamagmatites from Pan-African Orogen in Togo, West Africa*. ESI Preprints. <https://doi.org/10.19044/esipreprint.7.2023.p564>

Abstract

The Dahomeyide orogen in Togo and adjoining parts of SE-Ghana and Benin records the suture of West Africa Craton (WAC) into NW Gondwana. The suture zone corresponds to a narrow and lithologically diverse area with HP granulites complexes. In Togo, the Djabatoure massif, located in the central part of Togo, belongs to the suture zone. The aim of this paper is to present the petrographic and geochemical characteristics of the Djabatoure massif in order to understand its implication in the geodynamic evolution of the Dahomeyide belt in Togo. The methodology

implemented is based on a synthesis of previous works, a petrographic study of twenty thin sections and a geochemical study through discrimination diagrams of fifteen samples. The result shows that the Djabatoure massif is composed of granulites, pyroxenites, amphibolites, talcschists and gneisses. These rocks were equilibrated under granulite facies conditions and subsequently partially retrogressed to amphibolite facies. They display tholeiitic affinity, enriched LREE and negative anomalies in Nb, Zr and Ti indicating subduction zone magma. These features are consistent with protoliths of tholeiites, N-MORB and volcanic arc basalts affinities. The Djabatoure massif rocks were set up in oceanic environment and are thought to have been derivated from a metasomatized mantle.

Keywords: Petrography, Geochemistry, Djabatoure massif, Pan-African orogen, West Africa

1. Introduction

In the Pan-African Dahomeyide belt, the basic to ultrabasic complexes Derouvarou (Benin), Kabye-Kpaza, Djabatoure-Anie, Agou-Ahito (Togo), and Shaï or Akuse (Ghana) draw a submeridian mountainous belt materializing the suture zone (Ménot and Séddoh, 1980; Agbossoumondé, 1998; Attoh, 1998; Agbossoumondé et al., 2001; Tairou and Affaton, 2013). The Djabatoure massif located in the central part of Togo represents one of the most remarkable morphostructures of this belt. It is defined, like the other massifs of this suture zone, as a highly metamagmatic ophiolitic-type unit. It is composed of ultramafic, gabbroic metacumulates, metagabbros, metadolerites, metabasalts, pyroxenites, amphibolites and granulites (Ménot, 1977, 1980, 1982; Ménot and Seddoh, 1980; Agbossoumondé, 1998; Duclaux, 2003; Agbossoumondé et al, 2007; Kpanzou, 2017, 2023; Kpanzou et al., 2019, 2022). Moreover, the work of some authors (Duclaux, 2003; Sabi, 2007; Agbossoumondé et al., 2007) has partially addressed the petrographic and geochemical characteristics of all the massifs of the togolese segment of the suture zone. According to this work, the Djabatoure massif is composed of dolerites and gabbros and is formed by several episodes of magma injections into the migmatitic basement of the Touareg Shield. It shows a signature of continental tholeiite emplaced in the continental crust and having undergone a slow cooling during its exhumation. The rocks of Djabatoure massif have undergone metamorphism in the granulite facies with a retrogression in the amphibolite facies associated with the nappes setting. These are metaluminous rocks with a tholeiitic signature emplaced in a continental or oceanic arc context and exhumed during the Pan-African tangential phase.

Despite these efforts to clarify the petrography and geochemistry of the Djabatoure massif metamagmatites, detailed petrographic and geochemical characterization of the rocks of this massif remains a source of investigation. Therefore, the present study is a contribution to the detailed petrographic and geochemical study of the Djabatoure massif metamagmatites. It proposes to (i) determine the petrography of these rocks, (ii) classify them and identify their origin, (iii) and identify their geodynamic environment.

2. Geological context

The Pan-African Dahomeyide belt results from the collision between the eastern portion of the West African Craton (WAC) and the Benino-Nigerian basement and Touareg Shield set (Caby et al., 1981; Attoh et al., 1997; Ganade de Araujo et al., 2014a). It is subdivided into three structural zones which are from west to east: the external zone, the suture zone and the internal zone (Affaton, 1990) (Figure 1).

The external zone corresponds to the Buem and Atacora structural units, representing the metasedimentary units of the Volta Basin, the orthogneissic units of Kara-Niamtougou, and the granitoids of the Kpalimé-Amlamé pluton (Affaton, 1990; Affaton et al., 1991; Agbossoumondé et al., 2007; Tairou et al., 2009; Aidoo et al., 2020; Kwayisi et al., 2021; Tairou et al., 2022). The external zone overlaps the WAC and its sedimentary cover.

The suture zone is materialized by a submeridian alignment of complexes: Derouvarou in Benin, Kabye-Kpaza, Djabatoure-Anie and Agou-Ahito in Togo, and Akuse or Shaï in Ghana. These complexes are composed of mafic to ultramafic rocks with mainly granulite or sometimes eclogite facies, witnesses of the Pan-African crustal thickening (Ménot and Séddoh, 1985; Attoh, 1998; Agbossoumondé et al., 2001; Attoh and Morgan, 2004; Aidoo et al., 2020; Kpanzou, 2023).

The internal zone corresponds to the peneplain representing the remobilized Benin-Nigerian shield (BNS) in the Pan-African belt (Affaton, 1990). It is composed of gneisso-migmatitic, metasedimentary (schists, marbles and quartzites) and Pan-African granitoid units (Affaton et al., 1991; Caby and Boessé, 2001; Alayi, 2018).




Figure 1. Simplified geologic map showing the main structural domains of the Pan-African Dahomeyide belt and its foreland (Affaton, 1990; slightly modified) with the location of the Djabatoure massif.

3. Methodology

In this study, the methodological approach is based on a number of processes including documentation, field work, laboratory analysis, data processing and analysis. The documentation was used to synthesize previous work on the regional and/or local geology of the study area. The field campaign allowed us to take fifty samples and describe them in a macroscopic way. We georeferenced the outcrops and sampling stations

using a Garmin eTrex Legend H GPS. Twenty (20) thin sections were made to determine textural and mineralogical compositions. We selected fifteen (15) samples for whole-rock analysis performed at the Centre for Scientific Instrumentation of the University of Granada (CIC-UGR) in Spain. The analytical approach adopted is as follows:

- (i) Major elements oxides were determined with a Philips Magix Pro (Pw-2440) X-ray fluorescence (XRF) equipment after melting the sample in a solution with tetra lithium borate. The characteristic precision as determined from standards AN-G and BEN, was better than $\pm 1.5\%$ (relative error) for an analyte concentration of 10 wt.%. The iron content is expressed as FeO* total. The molar ratio MgO/(MgO+FeO*) is abbreviated Mg#. Zirconium was determined in the same machine using the same glass beads with a precision better than $\pm 0.2\%$ for 5 ppm Zr. Loss on Ignition (LOI) was determined by weight difference before and after ignition of samples in a furnace. In the diagrams, oxide concentrations are reported in an anhydrous (volatile free) basis.

- (ii) Trace elements, except Zr, were determined by an Inductively Coupled Plasma-Mass Spectrometry (ICP-MS) after HNO₃ + HF digestion of 0.1000 g of sample powder in a Teflon-lined vessel at 180°C and 200 psi for 30 min, evaporation to dryness and subsequent dissolution in 100 ml of 4 vol.% HNO₃; the precision, as determined from standards PMS, WSE, UBN, BEN, BR and AGV run as unknowns, was better than $\pm 2\%$ for analyte concentrations of 50 ppm and $\pm 5\%$ for analyte concentrations of 5 ppm.

The results of the chemical analyses are reported in Table 1. The location of the analyzed samples is shown in Figure 2. Field and chemical data were compiled in Excel and imported into different software (GCD kit, QGIS ...) for specific processing. For the geochemical data processing, we proceeded to the normalization of major elements to 100% under anhydrous basis.

4. Results

4.1. Petrography

The Djabatoure massif metamagmatites (Figure 2) are essentially composed of granulites (garnet-bearing or garnet-free), pyroxenites, amphibolites, talcschists and gneisses (garnet-bearing or garnet-free). Representative rocks samples were selected for petrographic studies to determine the textures and mineralogical compositions.




Figure 2. Schematic map of the Djabatoure massif showing the location of the samples analyzed (modified from the geological map of Sylvain et al., 1986).

1: micaschists and graphitic schists; 2: sericite- and chlorite-bearing schists; 3: feldspathic quartzites; 4: garnet-bearing gneiss; 5: 2-mica-bearing paragneiss; 6: biotite-bearing micaschists; 7 : 2-mica-bearing gneiss; 8: kyanite- and garnet-bearing gneiss; 9: biotite- and amphibole-bearing gneiss; 10: clear metagabbroic granulites; 11: 2-mica- and amphibole-bearing gneiss; 12: amphibolites; 13 : Metavolcanites and associated albitic schists; 14: tectonized feldspathic quartzites; 15: basic charnockitic orthogneisses; 16: kyanite-bearing quartzites; 17: crystalline dolomites (marbles); 18: dark metagabbroic granulites; 19: pyroxenites; 20: serpentinites; 21: quartz veins; 22: muscovite- and or sericite-bearing quartzites; 23: overlapping contact; 24: towns and villages; 25: samples.

4.1.1. Granulites

The granulites observed and described in the field are of two types: garnet-free granulites and garnet-bearing granulites.

4.1.1.1. Garnet-free granulites

They outcrop at Sotouboua, Niki-Niki and Yeloum massifs. They are cut into milky white to grayish banks and balls (Figure 3a). Microscopically, they show granoblastic to double coronite textures (Figure 3b). The coronites are evidence of re-equilibration reactions between plagioclase and pyroxene blasts. Reactions in which clinopyroxene develops in the crown around orthopyroxene and amphibole in that around plagioclase. These reactions of re-equilibration often take place in a context of pressure and rather high temperature. The essential minerals are plagioclase, clinopyroxene, orthopyroxene and amphibole. Plagioclases are more abundant and in phenocrysts with polysynthetic twin and more or less clear rolling extinction. Orthopyroxenes (hypersthene) are deformed and show numerous cleavages, microfractures undulate extinction. Clinopyroxenes (diopside) occur as slightly microfractured or intensely cracked subautomorphic blasts, either as small crystals in the internal crown around the orthopyroxene. Amphiboles in elongated subautomorphic phenocrysts contain oxide inclusions. They are generally very little represented and constitute either early minerals in the form of large patches associated with pyroxenes or acicular minerals forming crowns around orthopyroxenes. Quartz is quite rare and occurs in xenomorphic form in inclusion in plagioclase and amphibole.

4.1.1.2. Garnet-bearing granulites

They are observed at Sotouboua massif. They are homogeneous, with a foliation of alternating light quartzo-feldspathic and dark amphibole and pyroxene layers, to which are added numerous millimetric to centimetric garnets, reddish in veinlets and wire rods (Figure 3c). Microscopically, they have a granoblastic texture consisting of plagioclase, pyroxene (orthopyroxene), amphibole and garnet (Figure 3d). Plagioclase is abundant and exists as xenomorphic and sometimes automorphic phenocrysts with polysynthetic macles. Orthopyroxene (hypersthene) is subautomorphic with strong relief and nearly orthogonal cleavages. Amphibole (hornblende) is scarce and occurs in small, more or less elongated sections of bluish green to brown color. Garnet is subautomorphic to automorphic and sometimes globular. It is very abundant in ferromagnesian beds and contains inclusions of quartz, plagioclase, epidote and opaque minerals (ilmenite, rutile and magnetite).

4.1.2. Pyroxenites

They outcrop at Sotouboua, Niki-Niki and Yeloum massifs. They are often cumulates with coarse or medium grains of pyroxene (Figure 3e). They are dark gray to black, massive in structure and flow in very dense balls. Microscopically, they display a granoblastic texture made of jointed

pyroxenes, with green hornblende crystallized at their borders (Figure 3f). They are mainly composed of clinopyroxene, orthopyroxene and amphibole. Clinopyroxenes (diopside) are the most abundant. They are slightly colored in green and show numerous cracks and undulate extinction. Orthopyroxenes (hypersthene) are pinkish colored with undulate extinction. They are much cleaved and contain inclusions of opaque minerals (ilmenite, rutile and magnetite). Amphiboles (green hornblende) are elongated phenocrysts joined to pyroxenes. They also constitute small crystals forming a reaction crown around orthopyroxenes (result of the reaction between orthopyroxenes and clinopyroxenes).



Figure 3. Macroscopic (a, c and e) and microscopic (b, d and f) photography of the Djabatoure massif rocks. (a and b): coronite garnet-free granulite; (c and d): garnet-bearing granulite; (e and f): pyroxenite. Opx: orthopyroxene; Cpx: clinopyroxene; Pl: plagioclase; Hbl: hornblende; Grt: garnet.

4.1.3. Amphibolites

They are observed at Sotouboua and Yeloum massifs. They are dark grey rocks, with a foliated structure and fine grains that are cut into platelets. On the field, they are crossed by veinlets and quartzo-feldspathic wire rods (Figure 4a). Microscopically, they have a nematogranoblastic texture with amphibole, plagioclase and quartz minerals (Figure 4b). Amphibole is represented by green hornblende in elongated bluish-green or brownish elongated sections. Plagioclase is more or less abundant and present polysynthetic twin. Quartz, with rolling extinction, is not very abundant and is found in association with plagioclase in the clear beds.

4.1.4. Talcschists

They outcrop in balls at Niki-Niki massif. They are milky white to greenish, with a schistose or foliated structure (Figure 4c). Microscopically, they have a porphyroblastic texture with phenoblasts of clinopyroxene, talc and serpentine (Figure 4d). Clinopyroxene (diopside) occurs as intensely cracked phenocrysts. It sometimes recrystallizes into amphibole or serpentine and contain inclusions of opaque minerals (magnetite). Talc, often associated with chlorite is from retromorphosis of amphiboles.

4.1.5. Garnet-bearing gneiss

These are mostly amphibole- and garnet-bearing gneisses that outcrop around granulites. They are grayish, of foliated structure and with millimetric to centimetric garnets (Figure 4e). Microscopically, they show a granoblastic texture composed of feldspars, quartz, biotite, amphibole and garnet (Figure 4f). Feldspars are subautomorphic and polysynthetic twin. Quartz shows undulate extinction. Amphibole (green hornblende) forms elongated, cracked blasts with opaque mineral inclusions. Garnets in automorphic porphyroblasts are often cracked and contain inclusions of feldspars, quartz and oxides.




Figure 4. Macroscopic (a, c and e) and microscopic (b, d and f) photography of the Djabatoure massif rocks. (a and b): amphibolite; (c and d): talcschist; (e and f): garnet-bearing gneiss. Qtz: quartz; Pl: plagioclase; Cpx: clinopyroxene; Hbl: hornblende; Bt: biotite; Grt: garnet; Srp: serpentine; Tlc: talc; Op: opaque.

4.2. Geochemistry

4.2.1. Major elements distribution

The Djabatoure massif rocks are characterized by silica contents in the range of 40.26% to 54.14% (Table 1). They are mainly granulites, pyroxenites and amphibolites. The analyzed samples show high values of Al_2O_3 up to 21.34% and Sr up to 580.91 ppm, which implies their richness in plagioclase. MgO contents reach 19.14% and Fe_2O_3 up to 13.97% indicating pyroxene accumulation.

According to Harker diagrams (Figure 5): granulites show a positive correlation between SiO_2 vs TiO_2 , FeOt , Na_2O , K_2O , P_2O_5 and negative with Al_2O_3 , MgO and CaO ; pyroxenites show a positive correlation between SiO_2 vs MgO and negative with Al_2O_3 , TiO_2 , CaO , Na_2O , K_2O and P_2O_5 ; amphibolites show a positive correlation between SiO_2 vs Na_2O , K_2O , P_2O_5 and negative with Al_2O_3 , TiO_2 , FeOt and CaO .

Table 1. Results of geochemical analyses of major elements (%), traces elements (ppm) and rare earth elements (ppm) of the Djabatoure massif rocks.

Lithology	Granulites								Pyroxénites					Amphibolites	
Ech.	85B	89	90	105	115	118	119A	120	64A	64B	100B	116	145	65	119B
Xcood.	E1°1201,9	E1°1151,3	E1°1010,8	D7°5528,8	D7°5742,8	D7°5705,3	D7°5815,8	D7°5808,3	E1°0236,3	E1°0236,3	E1°1031,1	D7°5801,4	E1°0231,4	E1°0245,7	D7°5815,8
Ycood.	N8°2112,1	N8°2148,2	N8°2004,1	N8°3412,4	N8°3310,8	N8°3325,1	N8°3248,8	N8°3237,4	N8°0823,1	N8°0823,1	N8°2333,1	N8°3310,3	N8°0839,9	N8°0809,9	N8°3248,8
Majors (%)															
SiO ₂	50,64	51,86	49,42	54,14	49,70	50,34	47,80	48,73	47,05	51,86	49,10	47,76	52,36	50,33	40,26
Al ₂ O ₃	7,90	17,09	16,60	17,88	18,89	14,69	19,83	18,20	9,85	5,16	15,09	7,56	6,55	14,71	21,34
Fe ₂ O ₃	8,74	6,94	10,87	9,55	7,49	12,20	7,74	7,86	10,81	9,12	9,00	10,69	11,45	12,54	13,97
MnO	0,15	0,12	0,14	0,15	0,12	0,27	0,11	0,12	0,17	0,14	0,15	0,21	0,19	0,21	0,11
MgO	13,76	9,63	6,19	4,09	6,48	8,80	10,52	9,98	14,60	19,14	10,90	15,44	17,52	6,59	6,83
CaO	16,05	10,37	9,29	8,43	12,82	9,99	11,27	11,48	12,54	10,95	11,87	14,85	10,34	9,38	12,27
Na ₂ O	0,81	2,51	2,70	3,40	2,82	2,28	2,24	2,44	1,73	0,78	1,43	1,24	0,98	3,25	2,00
K ₂ O	0,16	0,20	0,88	0,49	0,17	0,17	0,04	0,07	0,36	0,10	0,33	0,11	0,05	0,23	0,11
TiO ₂	0,59	0,38	1,18	1,17	0,67	0,93	0,24	0,28	1,19	0,55	0,71	0,93	0,51	1,47	1,73
P ₂ O ₅	0,02	0,04	0,30	0,23	0,08	0,16	0,02	0,02	0,10	0,03	0,07	0,02	0,02	0,13	0,03
PAF	0,62	0,45	1,92	0,00	0,68	0,00	0,00	0,23	1,02	1,74	0,90	0,53	0,02	0,80	0,74
Total	99,44	99,59	99,49	99,53	99,92	99,83	99,81	99,41	99,42	99,57	99,55	99,34	99,99	99,64	99,39
Traces (ppm)															
Rb	4,89	3,29	19,61	3,37	1,96	1,56	0,14	0,44	2,23	0,67	5,52	0,34	0,57	4,15	0,28
Ba	42,72	96,26	240,35	215,04	53,30	82,83	20,91	77,80	54,42	49,18	83,92	36,66	67,11	49,98	26,93
Nb	1,14	1,15	2,67	5,26	1,80	3,87	0,84	0,76	2,66	1,24	2,26	1,30	1,24	2,50	1,38
Ta	0,08	0,09	0,16	0,37	0,15	1,49	0,28	0,07	0,15	0,05	0,14	0,08	0,47	0,19	0,15
Sr	166,39	418,23	580,91	438,62	355,62	298,91	300,27	284,90	113,94	55,10	377,78	117,89	130,85	162,12	514,87
Zr	35,50	23,00	90,20	80,00	35,60	35,60	13,60	14,40	39,50	24,80	51,00	23,10	24,00	81,60	27,50
Y	19,38	7,89	18,26	20,17	11,55	18,83	3,18	5,17	18,76	12,60	14,32	16,32	12,01	30,74	13,31
Hf	1,07	0,28	0,75	0,57	0,32	0,46	0,00	0,00	1,28	0,49	0,68	0,59	0,52	0,64	0,41
Ni	359,76	189,36	64,01	16,84	71,98	116,29	200,46	214,46	497,54	580,30	150,51	338,84	580,91	34,14	92,97
Cr	2269,25	363,63	123,63	33,90	270,92	440,88	317,29	355,17	949,85	1715,01	233,42	1842,86	939,15	15,49	18,09
U	248,01	113,08	304,38	226,53	136,56	164,86	49,45	74,89	298,24	195,28	173,92	187,78	205,31	314,93	452,41
Th	0,13	0,04	0,17	0,12	0,04	0,05	0,00	0,00	0,16	0,07	0,09	0,01	0,02	0,11	0,00
U	0,00	0,00	0,01	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00
Sc	56,84	22,83	26,48	24,54	29,01	41,09	17,17	25,04	50,07	46,48	39,15	47,41	40,57	41,16	27,83
Co	60,92	53,13	49,84	54,77	46,19	208,01	94,81	66,41	65,34	67,64	60,35	64,71	130,98	57,99	82,66
Cu	123,64	24,76	87,87	17,75	18,77	48,78	42,29	95,65	151,20	80,19	119,31	94,97	194,70	69,17	204,36
Zn	49,25	50,18	85,15	84,18	44,19	108,90	38,99	38,98	76,52	59,47	66,72	67,19	71,18	91,75	72,77
Mo	1,77	2,84	2,55	5,10	2,68	2,91	3,69	2,38	1,77	1,31	2,46	2,28	2,89	2,43	2,17
Cs	0,33	0,24	0,53	0,14	0,12	0,12	0,08	0,08	0,09	0,10	0,34	0,06	0,08	0,12	0,08
Li	6,91	5,42	8,72	8,23	2,64	2,98	0,00	0,57	2,79	1,08	9,47	0,43	1,44	6,39	0,79
Be	0,30	0,37	0,78	0,97	0,34	0,54	0,21	0,21	0,43	0,28	0,40	0,21	0,30	0,63	0,24
Ga	10,13	14,29	20,14	20,64	17,71	15,10	14,57	15,18	14,12	7,65	14,38	10,29	9,41	16,83	20,87
Sn	0,37	0,03	0,71	0,33	0,16	0,38	0,00	0,00	0,71	0,23	0,35	0,32	0,24	0,63	0,37
Tl	0,04	0,03	0,09	0,03	0,03	0,03	0,02	0,04	0,03	0,02	0,04	0,02	0,02	0,04	0,02
Pb	0,65	1,30	3,56	3,70	1,43	1,62	0,10	0,16	0,59	0,45	1,68	0,00	0,29	1,65	0,22
REE (ppm)															
La	4,98	3,64	10,67	10,52	3,40	5,65	0,72	0,68	7,23	3,54	4,99	3,03	2,48	5,24	2,43
Ce	12,86	7,44	25,12	23,70	8,34	13,06	1,86	1,83	18,68	8,95	11,75	7,91	6,49	13,52	6,30
Pr	1,95	1,01	3,51	3,20	1,21	1,90	0,27	0,30	2,83	1,34	1,68	1,30	1,11	1,92	1,02
Nd	9,57	4,70	16,24	14,86	6,12	9,58	1,41	1,65	14,58	6,92	8,27	7,62	5,95	10,08	5,55
Sm	3,00	1,22	3,89	3,71	1,77	2,80	0,39	0,55	3,95	2,03	2,31	2,74	1,84	3,30	2,07
Eu	0,93	0,38	1,29	1,23	0,50	0,85	0,12	0,20	1,28	0,58	0,81	0,77	0,63	1,08	0,67
Gd	2,48	1,19	2,94	2,98	1,46	2,33	0,42	0,64	3,06	1,55	2,39	2,30	1,99	3,78	1,87
Tb	0,46	0,21	0,46	0,50	0,26	0,43	0,07	0,11	0,51	0,29	0,36	0,40	0,33	0,68	0,34
Dy	2,96	1,27	2,85	3,09	1,75	2,82	0,57	0,76	3,04	1,92	2,18	2,61	1,81	4,42	2,19
Ho	0,64	0,28	0,60	0,66	0,40	0,62	0,14	0,20	0,63	0,43	0,48	0,55	0,41	0,99	0,46
Er	1,64	0,86	1,32	1,69	1,07	1,64	0,42	0,65	1,58	1,18	1,31	1,42	1,15	2,47	1,20
Tm	0,24	0,14	0,22	0,25	0,16	0,25	0,07	0,11	0,23	0,18	0,20	0,21	0,18	0,36	0,18
Yb	1,43	0,87	1,32	1,49	0,99	1,50	0,39	0,66	1,32	1,07	1,17	1,19	1,05	2,18	0,99
Lu	0,21	0,13	0,19	0,23	0,14	0,23	0,06	0,09	0,19	0,15	0,17	0,17	0,16	0,33	0,14
Eu/Eu*	1,02	0,94	1,14	1,11	0,93	1,00	0,89	1,01	1,10	0,98	1,03	0,92	0,99	0,92	1,02
(La)N	12,97	9,48	27,79	27,40	8,85	14,71	1,88	1,77	18,83	9,22	12,99	7,89	6,46	13,65	6,33
(Sm)N	12,95	5,27	16,80	16,02	7,64	12,09	1,68	2,37	17,06	8,77	9,97	11,83	7,95	14,25	8,94
(Gd)N	8,09	3,88	9,59	9,72	4,76	7,60	1,37	2,09	9,98	5,06	7,80	7,50	6,49	12,33	6,10
(Yb)N	6,22	3,78	5,74	6,48	4,30	6,52	1,70	2,87	5,74	4,65	5,09	5,17	4,57	9,48	4,30
(La/Sm)N	1,00	1,80	1,65	1,71	1,16	1,22	1,11	0,75	1,10	1,05	1,30	0,67	0,81	0,96	0,71
(Gd/Yb)N	1,30	1,03	1,67	1,50	1,11	1,17	0,81	0,73	1,74	1,09	1,53	1,45	1,42	1,30	1,42
(La/Yb)N	2,09	2,51	4,84	4,23	2,06	2,26	1,11	0,62	3,28	1,98	2,55	1,53	1,41	1,44	1,47
ΣREE	43,35	23,34	70,82	68,11	27,57	43,66	6,91	8,43	59,11	30,13	38,07	32,22	25,58	50,35	25,41




Figure 5. Oxides (TiO_2 , FeOt , Na_2O , K_2O , P_2O_5 , Al_2O_3 , MgO , CaO) vs SiO_2 variation diagrams in Djabatoure massif rocks (Harker, 1909).

4.2.2. Traces and rare earth elements distribution

In the trace elements (Rb, Sr, Nd, Ba, Nb, Y, Ni, Cr) versus Zr diagrams (Harker, 1909) (Figure 6): granulites show a positive correlation between Zr vs Rb, Sr, Nd, Ba, Nb, Y and negative with Ni and Cr; pyroxenites show a positive correlation between Zr vs Rb, Sr, Nd, Ba, Nb, Y and negative with Ni and Cr; amphibolites show a positive correlation between Zr vs Rb, Nd, Ba, Nb, Y and negative with Sr and Ni.

Rare earth spectra normalized to chondrites (Nakamura, 1974) (Figure 7a) show that the rocks of the Djabatoure massif are moderately

fractionated ($0.62 \leq (\text{La}/\text{Yb})_N \leq 4.84$). They are slightly rich in light rare earth elements ($0.67 \leq (\text{La}/\text{Sm})_N \leq 1.80$) and show generally flat heavy rare earth elements spectra ($0.73 \leq (\text{Gd}/\text{Yb})_N \leq 1.74$). These rocks record positive anomalies in Pr, Ho, Tm and negative in Ce, Nd, Gd, Dy, Er, Yb.

Multielement spectra normalized to the early mantle (McDonough and Sun, 1995) (Figure 7b) show that the rocks of the Djabatoure massif have positive anomalies in Ba, Ta, Sm, Y and negative in Rb, U, Nb, Pb, Zr.




Figure 6. Traces elements (Rb, Sr, Nd, Ba, Nb, Y, Ni, Cr) versus Zr variation diagrams in Djabatoure massif rocks (Harker, 1909).




Figure 7. (a) Rare earth elements spectra normalized to chondrites (Nakamura, 1974) and (b) trace elements spectra normalized to the primitive mantle (McDonough and Sun, 1995) of the Djabatoure massif rocks.

4.2.2. Type of the rocks and geodynamic context

The Djabatoure massif rocks are distributed in the TAS diagram of Middlemost (1994) (Figure 8a) in the gabbroic diorites and gabbros fields. The diagram of Irvine and Baragar (1971) (Figure 8b) indicates that the rocks of this massif have a tholeiitic affinity. The diagram of Shand (1943) (Figure 8c) shows that these rocks are metaluminous. The Meschede (1986) diagram (Figure 8d) shows that the rocks of the Djabatoure massif fall into two domains: N-MORB (D), and intraplate tholeiitic and volcanic arc basalts (C).




Figure 8. Geochemical classification plots of Djabatoure massif rocks. (a): $\text{Na}_2\text{O}+\text{K}_2\text{O}$ vs SiO_2 (Middlemost, 1994); (b): $\text{FeOt}-(\text{Na}_2\text{O}+\text{K}_2\text{O})-\text{MgO}$ (Irvine and Baragar, 1971); (c): A/NK vs A/CNK (Shand, 1943); (d): $\text{Zr}/4-\text{Nb}-\text{Y}$ (Meschede, 1986) [AI = intraplate alkaline basalts; AII = intraplate alkaline basalts and tholeiites; B = E-type MORB; C = intraplate tholeiites and volcanic arc basalts; D = N-type MORB].

5. Discussion

5.1. Petrography and metamorphic evolution

The petrographic study of the Djabatoure massif shows a diversity of rocks: granulites, pyroxenites, talcschists, amphibolites and gneisses. The different mineralogical associations of the samples described are presented in Table 2. They show the different metamorphic episodes that affected these rocks.

Coronitic granulites are generally garnet-free. Coronitic textures are the result of reactions between plagioclases and orthopyroxenes in which clinopyroxene develops in the crowns around orthopyroxenes and amphiboles in those around plagioclases. This amphibole crown is late and results from the hydration of clinopyroxenes.

Garnet-bearing granulites represent facies without coronites. They are characterized by the mineralogical assemblage of Pl+Opx+Hbl+Grt. Amphibolitization has led to the transformation of granulites (garnet-free or bearing-garnet) into amphibolites characterized by the Hbl+Pl+Qtz paragenesis, indicating the transition from granulite facies to amphibolite facies.

The pyroxenites have an Opx+Cpx+Pl+Hbl paragenesis and underwent a first retromorphosis in the amphibolite facies, with the replacement of pyroxenes by green hornblende, and a second one in the greenschist facies which led to serpentinization and talcschization.

The garnet gneisses present mineralogical associations with Pl+Qtz+Bt+Hbl+Grt witnessing a high-grade metamorphism with the presence of garnet in their composition.

Table 2. Mineralogical composition of the petrographic facies of the Djabatouré massif.

Petrographic facies	Main minerals	Accessory minerals
Garnet-free granulites	Pl+Cpx+Opx+Hbl	Ep+Opaque
Garnet-bearing granulites	Pl+Opx+Hbl+Grt	Opaque
Pyroxenites	Opx+Cpx+Pl+Hbl	Opaque
Amphibolites	Hbl+Pl+Qtz	Ep+Opaque
Talcschists	Cpx+Srp+Tlc	Ep+Opaque
Garnet-bearing gneiss	Pl+Qtz+Bt+Hbl+Grt	Opaque

5.2. Fractional crystallization

The correlations observed in the Harker variation diagrams of major and trace elements, the parallelism of rare earth elements spectra and multi-element spectra, indicate an evolution of the different facies from the unique magmatic source.

The slight negative anomalies Nb-Ta and the enrichment in Rb and Ba imply the influence of recycling of crustal material (crustal contamination). The positive anomaly in Sr suggests plagioclase accumulation. The good linear

correlation in the Harker diagrams reflects the intervention of a differentiation process by fractional crystallization in the evolution of the parent magmas of this massif rocks.

The increase of FeOt versus SiO₂ in the granulites indicates the accumulation of opaque minerals such as magnetite and ilmenite. The negative correlation of MgO vs SiO₂ in the granulites shows the fractionation of pyroxene and amphibole in these rocks. The increase of Na₂O and K₂O versus SiO₂ in granulites and amphibolites indicates their richness in acid plagioclase. The increase in P₂O₅ in granulites and amphibolites is indicative of either apatite crystallization or fluid circulation. The decrease of Al₂O₃ and CaO vs SiO₂ in granulites and amphibolites marks an early crystallization of calcium plagioclase in the least evolved terms. The negative correlation of MgO and CaO versus silica in granulites reflects the simultaneous fractionation of olivine, pyroxene and plagioclase in primary magmas.

The increase in Rb and Ba contents suggests the high fractionation of plagioclase in these rocks. The decrease in Ni and Cr contents vs Zr is consistent with the fractionation of ferromagnesian (pyroxene and amphibole) in these rocks. These elements could also be associated with MgO and FeOt in pyroxenes.

5.3. Source of magmas

The Djabatoure massif metamagmatites have an intermediate to ultrabasic composition. They have mineralogical and chemical characteristics that allow them to be classified as gabbros and gabbroic diorites according to the terminology of Middlemost (1994). The petrographic characteristics reflect the chemical composition of these rocks with the abundance of hornblende, the presence of orthopyroxene and clinopyroxene, as well as apatite as accessory mineral. Negative anomalies of elements such as niobium, phosphorus and titanium, observed on the multielement spectra are fingerprints commonly attributed to subduction zone magmas. The negative anomaly in Nb is characteristic of continental tholeiites and of crustal contamination or metasomatism from a mantle source.

The genesis of these rocks is therefore compatible with emplacement in a magmatic arc context with simultaneous crustal assimilation and fractional crystallisation of magma probably derived from continental crust.

5.4. Geotectonic environment

Based on the petrogeochemical data of the Djabatoure massif metamagmatites, such as the tholeiitic affinity rocks with metaluminous

signature, it can be proposed that they were emplaced in a probably extensive tectonic context, a context already mentioned by Duclaux (2003). Geotectonic discrimination diagrams indicate that these rocks were emplaced in various contexts: intraplate tholéiites, N-MORB and volcanic arc basalts. The negative anomaly Nb is characteristic of continental tholeiites and crustal contamination. It suggests a context of subduction zones in which Nb is trapped.

The Djabatoure massif rocks have similar geochemical characteristics to those of the Kabye and Agou massifs, such as high Sr and Ba values, a negative anomaly in Nb, and enrichment in light rare earth elements (LREE) compared to heavy rare earth elements (HREE). The rocks of this massif have been classified as typical late-orogenic magmas probably derived from a mantle source, enriched, metasomatized by plunging plate fluids and subducted sediments during a subduction event (Affaton, 1990; Agbossoumondé, 1998; Duclaux, 2003; Sabi, 2007) and subsequently contaminated by the continental crust.

Despite the lack of isotopic data, the analogies of petrogeochemical studies of the rocks of the Djabatoure massif with those of the Kabye and Agou massifs allow them to be classified in the same geodynamic context.

Conclusion

Petrographic studies of the Djabatoure massif have identified: granulites, amphibolites and gneisses representing foliated facies; massive to cumulative pyroxenites; schistose to laminated talcschists. Most of these rocks present mineralogies dominated by a primary paragenesis constituted by an abundance of plagioclase followed by clinopyroxene and a secondary paragenesis which is mainly represented by alteration minerals such as amphibole and iron and titanium oxides. These paragenesis show that these rocks underwent a metamorphism in the granulite facies with a retrogression in the amphibolite facies. Thus, the massif has undergone a metamorphic evolution from granulitization to retromorphosis in the amphibolite facies.

Geochemical characterization shows that the rocks of the Djabatoure massif display metaluminous and tholeiitic affinity. Inter-element variations indicate that these rocks evolved from a fractional crystallization process accompanied by crustal contamination. They were emplaced in various settings: intraplate tholéiites, N-MORB and volcanic arc basalts. They are crustal base rocks uplifted by Pan-African tectonics in crustal sets. Geochemical trends clearly indicate emplacement in a subduction context. The Djabatoure massif rocks show enrichment in LREE compared to HREE and sub-flat HREE spectra. The multi-element spectra show that most of the rocks are rich in mobile elements (Rb, Sr and Ba) and have negative

anomalies in Nb, P and Ti. These characteristics are similar to those of rocks from a mantle source enriched and metasomatized during a subduction event.

Acknowledgements

I would like to thank the authorities of the universities of the Coimbra Group who, through the "Coimbra Group Short Stay Scholarship Programme for young researchers from Sub-Saharan Africa", have granted me a laboratory research scholarship for the realization of geochemical analyses of rock samples. I also thank Professors Antonio GARCIA-CASCO and José María GONZÁLEZ-JIMÉNEZ, both professors-researchers at the Department of Petrology and Mineralogy of the University of Granada in Spain, who agreed to support my application for the scholarship and to carry out the chemical analyses of my rock samples in their laboratory. We acknowledge the anonymous reviewers for their constructive comments which improve the quality of the initial manuscript.

Conflict of Interests: The authors declare no conflict of interest.

References:

1. Affaton, P. (1990). Le bassin des Volta (Afrique de l'Ouest) : une marge passive d'âge protérozoïque supérieur, tectonisée au Panafricain (600±50 Ma). Edit. ORSTOM, Collection Etudes & Thèses, Paris, 500p. https://horizon.documentation.ird.fr/exl-doc/pleins_textes/pleins_textes_2/etudes_theses/31718.pdf
2. Affaton, P., Gelard, J.P., Simpara, N. (1991). Paléocontraintes enregistrées par la fracturation dans l'unité structurale de l'Atacora (Chaîne Panafricaine des Dahomeyides, Togo). C. R. Acad. Sci., Paris, t. 312 : 763-768.
<http://pascal-francis.inist.fr/vibad/index.php?action=getRecordDetail&idt=19679359>
3. Agbossoumondé, Y. (1998). Les complexes ultrabasiques de la chaîne panafricaine au Togo (Axe Agou – Atakpamé, Sud-Togo). Etude pétrographique, minéralogique et géochimique. Thèse Doct. Lab. Géol. Pétro. Univ. Jean Monnet St. Etienne Fr., 306p.
4. Agbossoumondé, Y., Ménot, R.-P., Guillot, S. (2001). Metamorphic evolution of Neoproterozoic eclogites from South Togo (West Africa). Jour. of Afr. Earth Sc. [https://doi.org/10.1016/S0899-5362\(01\)80061-0](https://doi.org/10.1016/S0899-5362(01)80061-0) 33, 227–244.
5. Agbossoumondé, Y., Ménot, R.-P., Paquette, J.L., Guillot, S., Yéssoufou, S., Perrache, C. (2007). Petrological and geochronological constraints on the origin of the Palimé–Amlamé

- granitoids (South Togo, West Africa): A segment of the West African Craton Paleoproterozoic margin reactivated during the Pan-African collision. *Gondwana Research* 12, 476-488. <https://doi.org/10.1016/j.gr.2007.01.004>
6. Aidoo, F., Sub, F.-Y., Liang, T., Nude, P.M. (2020). New insight into the Dahomeyide Belt of southeastern Ghana, West Africa: Evidence of arc-continental collision and Neoproterozoic crustal reworking. *Precambrian Research* 347 (2020) 105836. <https://doi.org/10.1016/j.precamres.2020.105836>.
 7. Alayi, G. (2018). Les granitoïdes tardifs de la chaîne panafricaine des Dahomeyides au Togo: étude pétro-structurale, géochimique et géochronologique. Thèse Doctorat, FDS, Univ. Lomé-Togo, 256p.
 8. Attoh, K. (1998). High-Pressure Granulite Facies metamorphism in the Pan-African Dahomeyide orogen, West Africa. *J. Geology*, 106 : 236-246. <https://doi.org/10.1086/516019>
 9. Attoh, K., Dallmeyer, R.D., Affaton, P. (1997). Chronology of nappe assembly in the Pan-Africa Dahomeyide Orogen, West Africa: evidence from ⁴⁰Ar/³⁹Ar mineral ages. *Precambrian research*, 82, pp. 153 – 171. [https://doi.org/10.1016/S0301-9268\(96\)00031-9](https://doi.org/10.1016/S0301-9268(96)00031-9)
 10. Attoh, K., Morgan, J. (2004). Geochemistry of high-pressure granulites from the Pan-African Dahomeyide orogen, West Africa: constraints on the origin and composition of lower crust. *Jour.of Afr.Earth Sci.*, vol. 39, pp. 201-208. <https://doi.org/10.1016/j.jafrearsci.2004.07.048>
 11. Caby, R., Bertrand, J.M., Black, R. (1981). Pan-African closure and continental collision in the Hoggar. Ifora segment, central Sahara. In Kröner A. (Eds) *Precambrian Plate Tectonics*. Elsevier, Amst, pp. 407-434. <https://www.scirp.org/%28S%28351jmbntvnsjt1aadkozje%29%29/reference/referencespapers.aspx?referenceid=1160492>
 12. Caby, R., Boessé, J.M. (2001). Pan–African nappe system in southwest Nigeria: the Ife– Ilesha schist belt. *Jour.of Afr.Earth Sci.*, vol. 33 n°2, pp. 211- 225. DOI : 10.1016/S0899-5362(01)80060-9
 13. Duclaux, G. (2003). Etude pétrologique et structurale des massifs basiques et ultrabasiques de la zone de suture panafricaine de la chaîne des Dahomeyides au Togo : Implications géodynamiques. *Mém. DEA, Lab. Dyn. Lithos. Univ. J. Monnet, St-Etienne*, 29p. <https://doi.org/10.13140/RG.2.2.28332.92809>
 14. Ganade de Araujo, C.E., Rubatto, D., Hermann, J., Cordani, U.G., Caby, R., Basei, M.A.S. (2014a). Ediacaran 2,500-km-long synchronous deep continental subduction in the West Gondwana

- Orogen. *Nature Communications*, 5:5198. doi: 10.1038/ncomms6198.
15. Harker, A. (1909). *The natural history of igneous Rocks*. Methuen and Co., London, 384p. <https://doi.org/10.1017/CBO9780511920424>
 - Irvine, T.N., Baragar, W.R.A. (1971). A guide to chemical classification of the common volcanic rocks. *Can. J. Earth Sci.*, 8: 523-548. <https://doi.org/10.1139/e71-055>
 16. Kpanzou, S.A.M. (2017). Etude pétrographique et structurale du massif de Djabatouré et des massifs adjacents. *Mém. Master, FDS, Univ. Lomé*, 31p.
 17. Kpanzou, S.A.M. (2023). Contribution à l'étude du complexe basique-ultrabasique de Djabatouré-Anié (Centre-Togo) : Caractéristiques pétrostructurales, géochimiques et indices de minéralisations associés. *Thèse de Doctorat unique, FDS, Univ. Lomé*, 232p.
 18. Kpanzou, S.A.M., Agbossoumondé, Y., Tairou, M.S. (2019). Caractérisation pétrostructurale du massif granulitique de Djabatouré. *J. Rech. Sci. Univ. Lomé (Togo), Spécial 2019*, 21(4-2) : 509-519. <https://www.ajol.info/index.php/jrsul/article/view/206992>
 19. Kpanzou, S.A.M., Agbossoumondé, Y., González Jiménez, J.M., Tairou, M.S., Garcia-Casco, A. (2022). Pétrologie et métallogénie des indices de Ni-Cr associés au massif basique-ultrabasique de Oké, Togo. *Afrique SCIENCE* 21(1) : 53-69. <https://www.afriquescience.net/PDF/21/1/5.pdf>
 20. Kwayisi, D., Elburg, M., Lehmann, J. (2021). Preserved ancient oceanic lithosphere within the Buem structural unit at the eastern margin of the west African craton, *LITHOS* (2021), <https://doi.org/10.1016/j.lithos.2021.106585>
 21. McDonough, W.F., Sun, S.S. (1995). The composition of the Earth. *Chemical Geology* 120, pp. 223–253. [https://doi.org/10.1016/0009-2541\(94\)00140-4](https://doi.org/10.1016/0009-2541(94)00140-4)
 22. Ménot, R.-P. (1977). Les massifs basiques et ultrabasiques antémétamorphiques de la bordure Ouest du mole Dahoméo-nigérian. *Essai de synthèse bibliographique. Ann. Univ. Bénin, Lomé*, pp. 53-94. <https://www.researchgate.net/publication/327822999>
 23. Ménot, R.-P. (1980). Les massifs basiques et ultrabasiques de la zone mobile panafricaine au Ghana, Togo et au Bénin. *Etat de la question. Bull. soc. Géol. Fr.*, 7 : 297-303. <https://doi.org/10.2113/gssgfbull.S7-XXII.3.297>
 24. Ménot, R.-P. (1982). Les éclogites des Monts Lato : un témoin de l'évolution tectono-métamorphique de la chaîne pan-africaine du

- Togo (Afrique de l'Ouest). *Ist. intern. Eclogite Conf., Terra Cognita*, 2, (3), 320p.
https://www.persee.fr/doc/geoly_0750-6635_1982_num_87_1_1513
- Ménot, R.-P., Seddoh, K.F. (1980). Le massif basique stratifié précambrien de Djabatoure-Soutouboua (région centrale du Togo, Afrique de l'Ouest). *Pétrologie et évolution métamorphique. Bulletin du B.R.G.M.* 4(4) : 319–337.
<https://www.researchgate.net/publication/260887781>
25. Ménot, R.-P., Seddoh, K.F. (1985). The eclogites of Lato Hills (South Togo, West Africa): relics from early tectonometamorphic evolution of the Pan-African orogeny. *Chemical Geology*, 50, pp. 313–330. [https://doi.org/10.1016/0009-2541\(85\)90126-3](https://doi.org/10.1016/0009-2541(85)90126-3)
26. Meschede, M. (1986). A method of discrimination between different types of mid-ocean ridge basalts and continental tholeiites with Nb-Zr-Y diagram. *Chemical Geology*, 56 : 207-218. [https://doi.org/10.1016/0009-2541\(86\)90004-5](https://doi.org/10.1016/0009-2541(86)90004-5)
27. Middlemost, E.A.K. (1994). Naming materials in the magma/igneous rock system. *Earth Science Reviews*, 37 (3-4), pp. 215-224. [https://doi.org/10.1016/0012-8252\(94\)90029-9](https://doi.org/10.1016/0012-8252(94)90029-9)
28. Nakamura, N. (1974). Determination of REE, Fe, Mg, Na and K in Carbonaceous and ordinary Chondrites. *Geochemica and Cosmochimica Acta*, 38, 757-775.
[http://dx.doi.org/10.1016/0016-7037\(74\)90149-5](http://dx.doi.org/10.1016/0016-7037(74)90149-5)
29. Sabi, B.E. (2007). Etude pétrologique et structurale du Massif Kabyè, Nord-Togo. Thèse Doctorat, FDS, Univ. Lomé, 256p.
30. Shand, S.J. (1943). *Eruptive Rocks. Their Genesis, Composition, Classification and Their Relations to Ore deposits.* 2nd edition, Murby London, 444p.
[https://www.scirp.org/\(S\(czeh2tfqyw2orz553k1w0r45\)\)/reference/ReferencesPapers.aspx?ReferenceID=1945539](https://www.scirp.org/(S(czeh2tfqyw2orz553k1w0r45))/reference/ReferencesPapers.aspx?ReferenceID=1945539)
31. Sylvain, J.P., Collart, J., Aregba, A., Godonou, S. (1986). Notice explicative de la carte géologique 1/500.000è du Togo, Mém. n°6, D.G.M.G./B.N.R.M., Lomé – Togo.
<https://scholar.google.com/scholar?q=+author:J.P.%20Sylvain>
32. Tairou, M.S., Affaton, P., Sabi, B.E., Seddoh, K.F. (2009). Tectono-metamorphic evolution of the Mo and Kara-Niamtougou Orthogneic Suites, Northern Togo. *Global Jour. of Geological Sciences*, vol. 7, n°2, pp. 93-100.
<https://www.researchgate.net/publication/287411921>
33. Tairou, M.S., Affaton, P. (2013). Structural Organization and Tectono-Metamorphic Evolution of the Pan-African Suture Zone: Case of the Kabye and Kpaza Massifs in the Dahomeyide Orogen in

- Northern Togo (West Africa). *International Journal of Geosciences* 04, 166-182. <https://doi.org/10.4236/ijg.2013.41015>
34. Tairou, M.S., Miningou, Y.M.W., Da Costa, Y.D., Kwekam, M. (2022). Petrostructural and Geochemical Characteristics of the Metamagmatites in the External Zone of the Dahomeyides Belt: Case of the Kantè Serpentinities (Northern Togo). *International Journal of Geosciences*, 2022, 13, 779-792. <https://www.scirp.org/journal/ijg>