



Assessing the Relationship between Short Birth-to-Pregnancy Interval and the Maternal and Perinatal Outcomes among Multiparous Women in Northern Ghana

Gilbert G. Agulu, BSc.PH, MPH, PhD (candidate)

Ghana Health Service, Regional Health Directorate
Private Mail Bag Bolgatanga, Upper East Region (UER), Ghana

Emmanuel K. Dzotsi, MD, MPH, MGCPs, FGCP

Ghana Health Service, Regional Health Directorate
Private Mail Bag Bolgatanga, UER, Ghana

Kofi Issah, MD, MPH

Family Health Division, Headquarters.

Ghana Health Service, Greater Accra, Ghana

Josephat A. Nyuzaghl, MGCPs, MPH, MD, BSc

Alemna D. Adogboba, BSc. MPH

Vincent A. Adongo, RN, BSc., MSc

Ghana Health Service, Regional Health Directorate
Private Mail Bag Bolgatanga, UER, Ghana

Doi: [10.19044/esipreprint.1.2024.p242](https://doi.org/10.19044/esipreprint.1.2024.p242)

Approved: 08 January 2024

Posted: 10 January 2024

Copyright 2024 Author(s)

Under Creative Commons CC-BY 4.0

OPEN ACCESS

Cite As:

Agulu G.G., Dzotsi E.K., Issah K., Nyuzaghl J.A., Adogboba A.D. & Adongo V.A.(2024). *Assessing the Relationship between Short Birth-to-Pregnancy Interval and the Maternal and Perinatal Outcomes among Multiparous Women in Northern Ghana*. ESI Preprints.

<https://doi.org/10.19044/esipreprint.1.2024.p242>

Abstract

Introduction: The World Health Organization (WHO) recommends an optimal Birth-to-Pregnancy Interval (BPI) of 24-59 months, or a 33-month interval between two successive births to reduce the risk of untoward maternal and newborn outcomes. Short Birth to Pregnancy Interval (SBPI), and unmet need for Family Planning (FP) are the major contributors to rapid population growth and increased maternal and newborn mortalities. The purpose of this study was to assess the adverse perinatal and maternal outcomes associated with SBPI among multiparous women in three municipalities of the Upper East Region (UER) of Ghana. Materials and

Methods:The study employed the Cross-sectional Design conducted among 904 women aged 15-49 years attending Antenatal Care (ANC) clinics in three municipalities in the UER who had at least two successive live births prior to data collection. Data was collected using structured questionnaire, incorporated into the electronic data collection tool, Kobo collect and administered by trained research assistants. Birth interval was categorized according to the WHO's classification, <24 months as SBPI, 24-≥59 months as Optimal Birth to Pregnancy Interval (OBPI) and >59 months as Long Birth to Pregnancy Interval (LBPI). **Results:** Of the 904 respondents, the majority (56.2%) had an OBPI, while 36.9% had a SBPI. Factors that influenced SBPI were parity, mode of delivery, and the educational status of women's partner. Participants with a higher parity (≥5 children) had 0.67 times the potential of spacing their births (AOR 0.67; 95% CI 0.46-0.98; p = 0.040). Women who experienced a Caesarean Section (CS) delivery were 3.28 times more likely to have LBPI (AOR 3.28; 95% CI 1.02-10.62; p=0.047). Respondents whose partners had secondary education had a 1.87 chance (AOR 2.07; 95% CI 1.09-3.96; p=0.027) of spacing their births. The birth complications reported were retained products of conception (41.9%), pregnancy-induced hypertension (27.9%), postpartum hemorrhage (11.6%), obstructed labor (10.5%), sepsis (38.1%), neonatal jaundice (23.8%), low-birth-weight (19%), and preterm birth (14.3%). **Conclusion and Global Health Implications:** The study recommends health professionals in direct contact with maternal, reproductive, and child health to be proactive in FP counseling, and to support women decision making.

Keywords: Birth-to-pregnancy interval; hemorrhage; maternal; multiparous; perinatal; postpartum; puerperal

1.0 Introduction

1.1 *Background of the Study*

Birth-to-Pregnancy Interval (BPI) which is also known as interbirth interval refers to the interval between the date of a live birth and the start of the subsequent pregnancy (Merdad & Ali, 2018). This is the period within which women recuperate from pregnancy and childbirth experience. It serves as a “fallow period” for physical, physiological, and psychological restoration after birth.

The World Health Organization (WHO) recommends an optimal BPI of 24–59 months, or a 33-month interval between two successive births, to reduce the danger of maternal, newborn, and perinatal deaths (Nausheen et al., 2021). This recommendation came preceding surveys highlighting the inapt effects of both short and long BPIs, such as low birth weight, preterm birth, small-for-gestational-age birth, and a high admission rate at the

neonatal intensive care unit (Bauserman et al., 2020; Merdad & Ali, 2018). Women with SBPIs are exposed to negative health consequences, especially when they have less time to physically convalesce from their previous pregnancy (Hailu & Gulte, 2016). Again, too-closed pregnancies increase the depletion of maternal nutrition, leading to maternal malnutrition (Mehra et al., 2019). A BPI of less than six months increases the probability of maternal mortality by 150% (95% CI 22% to 438%) and is linked with premature rupture of membrane, third trimester bleeding, postpartum endometriosis, and anemia (Nausheen et al., 2021). Recent study findings forecast that SBPI has the potential to cause autism in second-born children, particularly for pregnancies spaced less than 12 months apart (Conde-Agudelo et al., 2016). Another glaring consequence of SBI is, pregnant women are less likely to access Antenatal care (ANC) services due to the demand for care for the younger infants. For instance, babies are likely to be deprived of or compete for basic needs such as food, breastmilk, and parental attention (Molitoris, 2018). LBPIs, on the other hand, have been associated with an increased risk of pre-eclampsia (Nausheen et al., 2021). In sub-Saharan Africa (SSA), a survey in eight selected countries disclosed that most mothers ideally space their births. Nonetheless, SBPIs were reported in Chad (30.2%) and the Democratic Republic of the Congo (27.1%), while LBPIs were dominant in Southern and Eastern Africa, with Zimbabwe recording the highest percentage (Ajayi & Somefun, 2020). A recent study (Alhassan et al., 2022) revealed that Ghana has a SBPI prevalence of 49.7%, a low contraceptive prevalence rate, and high unmet needs for FP.

Several factors determine BPI; however, key among them include socio-demographic, biological, and sociological factors (Molitoris, 2018). Socio-demographic determinants like age, parity, type of residence, wealth index, education, and religion influence the BPI (Tesema et al., 2021). Studies show that prolonged postpartum amenorrhea is a primary biological factor in the long birth interval. According to Ajayi et al., contraceptive usage, length of breastfeeding, the death of the previous child, gender of the newborn, infrequent sexual intimacy, and the husband's occupation are strong sociological predictors of the BPI. These determinants were also common in other studies (Molitoris, 2018; Dehesh et al., 2020; Ayane et al., 2019).

Globally, SBPI and the unmet need for FP are the major contributors to rapid population growth. Data from the United Nations (UN), states that, the world is currently battling with a population of 7.6 billion people, which is projected to increase to 8.6 billion in 2030, 9.8 billion, and 11.2 billion in 2050 and 2100, respectively. Persistently elevated fertility rates due to SBPI pose health hazards for women and children, destroy human capital

investment, decline economic growth, and deplete environmental resources. Close to 4.7 million mothers, newborns, and children die annually in SSA, with a projected 265,000 mothers dying due to complications of pregnancy and childbirth (Tessema et al., 2020). Similarly, the population of Ghana is presently growing at a rate of 2.15% annually, though a gradual decline from a growth rate of 2.95% in 1985. The major indicators of the growth are deeply rooted in the high fertility rate of 3.89 births per woman and the country's mandates to reduce birth mortalities. Ghana has a high unmet need (30%) in FP (Ghana Maternal Health Survey, 2017). Ghana's maternal mortality ratio (MMR) is still unacceptably high (308 per 100,000 live births) as of 2017. In 2019, the neonatal mortality and infant mortality rates were 23.1 deaths per 1,000 live births and 33.9 deaths per 1,000 live births, respectively. The upper East Region (UER), for the past three years, has consistently recorded increasing MMRs. The region recorded 32 deaths (80.8/100,000LBs), 36 deaths (91.8/100,000LBs), and 43 deaths (98.1/100,000LBs) in 2019, 2020, and 2021 respectively (Ghana Maternal Health Survey, 2017). The situation is likely to further exacerbate if rapid targeted maternal interventions are not put in place.

A reduction in the unmet need for FP will minimize the high levels of unintended pregnancies, unsafe abortions, and maternal and neonatal mortality in the country (Guure et al., 2019). Again, by ensuring that women have access to comprehensive sexual and reproductive health services, including contraceptive usage, they will avoid becoming pregnant too early, too late, or too frequently and reduce avoidable pregnancy and childbirth-related deaths globally (Stover & Winfrey, 2017).

Despite the benefits associated with optimal birth spacing, the maternal and fetal outcomes in the study settings as well as in northern Ghana have not been explored. Again, the current BPI prevalence in the study sites remains unknown. Therefore, this study seeks to address these gaps by assessing the adverse perinatal and maternal outcomes associated with SBPI among multiparous women in three municipalities of the UER of Ghana.

1.1 Objectives of the Study

To describe the association between SBPI and adverse maternal and perinatal outcomes among multiparous women in the Kassena-Nankana Municipal (KNM), Bolgatanga Municipal (BoM), and Bawku Municipal (BaM) in the UER.

1.2 Specific Objectives

1. To estimate the proportion of short, optimal, and long BPIs among multiparous women in the three municipalities.

2. To determine the adverse effect of SBPI among multiparous women in the three municipalities.
3. To describe the predictors of BPI in the three municipalities.

1. Methods

2.1 Study Settings

The UER is one of the poorest regions of Ghana in terms of health and socioeconomic resources. The region is located in the north-eastern part of the country, with an estimated population of 1,318,523 people, with females constituting 48.4% of the population (Ghana Maternal Health Survey, 2017; GSS, 2017). The region shares borders with the Republic of Burkina Faso to the north, Togo to the east, the North-East Region of Ghana to the south, and the Upper West Region of Ghana to the west. Administratively, it is made up of 15 districts and municipalities and 101 sub-districts with health infrastructure such as a regional hospital, eight municipal and district hospitals, a few maternity homes and private faith-based facilities, health centers and multiple Community Health Planning and Services (CHPS) facilities. The study was conducted in three major municipalities in the region, namely KNM, BoM, and BaM.

2.2 Study Design

The study employed a cross-sectional design.

2.3 Study Population

All multiparous women who experienced at least two successive deliveries within the last five years preceding data collection.

2.4 Inclusion Criteria

All multiparous women in the age range of 15–49 years who had two successive live births within the last five years and stayed in the study area for six months prior to the data collection.

2.5 Exclusion Criteria

All nulliparous and uniparous women. Any woman below 15 years of age or above 49 years of age. Women whose children were severely sick and needed urgent treatment.

2.6 Data collection methods and instruments

Data was collected using a structured questionnaire that was adopted and modified from different literature⁴. Data collection was carried out by trained research assistants (four nurses and three midwives) and supervised by the investigators. The enumerators visited each selected health facility to

interview eligible mothers during ANC clinic days. The data collection was carried from October to December 2022.

2.7 Sample size determination

The sample size was calculated based on the average skilled deliveries (4,489) in the three municipalities in the last five years (2018-2022). The Cochran formula was used to determine sample size, $n = \frac{z^2 \times p(1-p)}{d^2}$, where n= required sample size, z= reliability coefficient (z-score) of 1.96 at 95% confidence level, p= estimated proportion of skilled delivery (50%); and d= margin of error of 5% (0.05). Since the prevalence of SBI was unknown in the region, we assumed 50%, this gives a sample of 384 respondents for each municipality.

2.8 Sampling Strategy

A multistage cluster sampling method was employed to recruit respondents for this study. It was conducted in four steps. **Step 1:** The study population was divided into clusters, of which the three municipalities serve as naturally occurring clusters. Within each cluster are the sub-municipalities. All sub-municipalities were considered for the sampling. **Step 2:** We randomly selected within each sub-municipalities' health facilities providing ANC service, of which there were a total of 67 health facilities (sample frame) in the three municipalities. **Step 3:** Out of the sample frame (67) we purposefully selected 46 health facilities. This included all health centers, hospitals, and CHPs facilities. **Step 4:** Of the 46 health facilities, 25 respondents were randomly selected using the systematic random selection method. In selecting the respondents, a proportionate quota was assigned to each facility depending on the population of ANC registrants. A total of 1,152 respondents were selected from the three municipalities for data collection.

2.9 Study Variables

The main exposure variable of interest was SBPI. A SBPI was defined in this study as an interval below 24 months occurring between the index and the current pregnancy. The major control variables included the age of the woman at first birth, sex of the index child, death of the index child, maternal and partner educational levels, place of residence, contraceptive usage, history of maternal infertility, etc. The main outcome variables were risks of maternal severe morbidity or a maternal near miss (e.g., mechanical ventilation, blood transfusion >2 units, admission into an intensive care unit, organ failure, etc.) and fetal outcomes such as stillbirth,

severe illness following delivery, delivery <28 weeks, and spontaneous and indicated preterm delivery (Schummers et al., 2018).

2.10 Statistical Analysis

Data analysis was done using descriptive statistics and Chi-square tests. In the descriptive analysis, frequency distribution and percentages were conducted to describe the demographic, socio-economic, obstetric, and childcare factors. In the Chi-square test, bivariate analysis was performed to identify factors associated with BPI. A multivariable logistic regression analysis was done to verify the variables linked with maternal and perinatal outcomes. The results of the regression analysis were presented using the Crude Odds Ratio (COR) and Adjusted Odds Ratio (AOR) with their corresponding Confidence Interval (CIs) and p-value. A p-value of <0.05 was considered statistically significant, showing the level of precision.

2.11 Ethical Approval/Informed Consent

Ethical approval was given by the Navrongo Health Research Centre Institutional Review Board of Ghana and the Ghana Health Service through the Upper East Regional Health Directorate. Written permission to carry out the study was sought from the municipal health authorities. Informed consent was written and explained to interviewees in their preferred languages. Participation was voluntary, and all participants had the right to withdraw from the study at any time. The identities of respondents were anonymized, and their records were made available to only the researchers. Participants' records were kept safely in a cabinet and locked. Literate participants were made to sign a copy of the consent form, while non-literate participants thumbed it. Consent was sought from legal guardians for participants who were under the age of 18.

a. Results

The study had a 21.5% dropout rate due to enumerators being unable to interview some respondents at the BaM study site as a result of an outbreak of conflict during the data collection period. Hence, out of the 1,152 respondents targeted, 904 were interviewed.

3.1 Sociodemographic Characteristics

In terms of educational attainment, 21.4% (193/904) of the respondents did not achieve any formal education, 56.6% (512/904) had primary and junior level education, and 22% (199/904) had secondary and tertiary education. More than a quarter (26.6%, 240/904) of the women partners did not have any formal education, while 31% (280/904) of them had secondary and tertiary education. For age categorization, only 16 (1.8%)

of the respondents were teenagers, whilst 63.2% (571/904) were in the age group of 25–34 years. Nearly 98% (883/904) of the women were married, with more than half of them (56%, 502/904) marrying at the age of 15–19 years, and 35% (314/904) marrying within the age brackets of 20–24 years. The majority, 92% (830/904) of the couples stay together. With regards to religion, 78.8% (712/904) of the women were Christians, and 18% (163/904) were Muslims. On ethnicity, about 42% (376/904) of the women were Frafra and 22%, (198/904) being Kassenas. As shown in Table 1 below.

Table 1. Demographic characteristics of respondents

Variable and category	Frequency	Percentage (%)
Municipal(n=904)		
BaM	135	14.9
BoM	384	42.5
KNM	385	42.6
Educational level of woman(n=904)		
No education	193	21.4
Primary	294	32.5
JHS	218	24.1
Secondary	126	13.9
Higher	73	8.1
Educational level of woman's partner (n=904)		
No education	240	26.6
Primary	189	20.9
JHS	195	21.6
Secondary	162	17.9
Higher	118	13.1
Age of woman (n=904)		
15–19	16	1.8
20–24	122	13.5
25–29	281	31.1
30–34	290	32.1
34–39	129	14.3
40–44	59	6.5
45–49	7	0.8
Mean \pm SD: 30.3 \pm 5.6		
Marital status(n=904)		
Married	883	97.7
Single	8	0.9
Separated	8	0.9
Divorced	1	0.1
Widow	4	0.4
Age of woman at marriage, n=896		
<15years	7	0.8
15–19years	502	56.0
20–24years	314	35.0
25–29years	61	6.8
30–34years	11	1.2
40–44years	1	0.1

Number of people in a household(n=904)		
<5 people	89	9.9
5–6 people	304	33.6
7 people and above	511	56.5
Religious affiliation(n=904)		
Christian	712	78.8
Islam	163	18.0
Traditionalist	18	2.0
No religion	11	1.2
Ethnic background (n=904)		
Frafra	376	41.6
Kassena	198	21.9
Nankana	154	17.0
Kusaasi	80	8.9
Builsa	26	2.9
Moshie	25	2.8
Bisa	13	1.4
Akan	10	1.1
Others ¹	22	2.4
Couple staying together? (n=904)		
No	74	8.2
Yes	830	91.8

¹ Hausa =4, Zambarama =3, Sissaala =3, Busanga =3, Fulami =3, Kantosis =2, Dagomba =2, Gonja =1, Guan =1

3.2 Association between delivery and newborn care, and BPI

As indicated in Table 2 below, the reason for preferred sex of the child was found to have a strong association with interbirth spacing ($\chi^2=15.1948$, $p=0.010$). Also, mode of delivery of the last child was a determinant of interbirth spacing ($\chi^2=12.9219$, $p=0.005$). We observed that women who delivered through CS spaced their births longer within 24–59 months. Again, parity ($\chi^2=5.1544$, $p=0.023$) was associated with BPI, as women with fewer than 1-3 children had SBPI compared to those with 4–7 children (64.3% versus 54.7%). Furthermore, the number of children a woman wishes to have ($\chi^2=14.1677$, $p<0.001$) and gravida ($\chi^2=15.4730$, $p<0.001$) have been found to be strongly linked with BPI. The period of commencement of sexual intercourse after delivery ($\chi^2=42.8910$, $p<0.001$) and those who received a postpartum FP method ($\chi^2=10.4212$, $p<0.001$) were strong predictors of BPI. The study found that women who breastfed their previous children ($\chi^2=5.8653$, $p=0.015$), practiced exclusive breastfeeding ($\chi^2=31.0284$, $p<0.001$), and the duration of exclusive breastfeeding ($\chi^2=45.3461$, $p<0.001$) were all strongly associated with BPI. Women who ever had a problem conceiving ($\chi^2=10.8256$, $p<0.001$) and the frequency of couples' sexual intercourse ($\chi^2=26.5771$, $p<0.001$) were found to be strong determinants of BPI.

Table 2. Association between delivery and newborn care and BPI

Variable	BPI		Total, n (%)	χ^2 (p- value)
	1-23 months, n (%)	24- \geq 59 months, n (%)		
Place of birth of last baby				6.8880 (0.229)
CHPS Compound	76 (31.1)	168 (68.9)	244	
Health Centre	67 (36.0)	119 (64.0)	(100.0)	
Clinic	25 (40.3)	37 (59.7)	186	
Hospital	161 (40.0)	242 (60.0)	(100.0)	
Home/TBA	2 (50.0)	2 (50.0)	62 (100.0)	
Aborted	3 (60.0)	2 (40.0)	403 (100.0)	
			4 (100.0)	
			5 (100.0)	
Sex of the index child				0.4322 (0.511)
Boy	195 (37.9)	320 (62.1)	515	
Girl	139 (35.7)	250 (64.3)	(100.0)	
			389 (100.0)	
Reasons for preferred sex of child				15.1948 (0.010) *
I have no preferred sex.	268 (36.8)	461 (63.2)	729	
Already had a girl (s).	7 (29.2)	17 (70.8)	(100.0)	
Already had a boy (s).	7 (35.0)	13 (65.0)	24 (100.0)	
My husband's preferred choice	19 (54.3)	16 (45.7)	20 (100.0)	
Child was my preferred sex.	33 (41.3)	47 (58.8)	35 (100.0)	
help in household activities	0 (0.0)	16 (100.0)	80 (100.0)	
			16 (100.0)	
Mode of delivery of last child				12.9219 (0.005) *
Spontaneous Vagina delivery (SVD)	321 (38.5)	512 (61.5)	833	
	7 (14.3)	42 (85.7)	(100.0)	
Caesarean section (CS)	0 (0.0)	1 (100.0)	49 (100.0)	
Vacuum extraction	6 (28.6)	15 (71.4)	1 (100.0)	
Forceps delivery			21 (100.0)	
Ever experienced birth complications during delivery (woman)?				1.8627 (0.394)
Yes	27 (31.4)	59 (68.6)	86 (100.0)	
No	307 (37.6)	510 (62.4)	817	
Do not remember	0 (0.0)	1 (100.0)	(100.0)	
			1 (100.0)	
Parity				5.1544 (0.023) *
1-3	234 (35.7)	422 (64.3)	656	
4-7	73 (45.3)	88 (54.7)	(100.0)	
			161 (100.0)	
Ever lost a child?				0.5106 (0.475)
Yes	49 (39.8)	74 (60.2)	123	
No	285 (36.5)	496 (63.5)	(100.0)	
			781 (100.0)	

Number of children ever lost				0.210
1	40 (39.6)	61 (60.4)	101	(0.900)
2	6 (37.5)	10 (62.5)	(100.0)	
3	2 (50.0)	2 (50.0)	16 (100.0)	
			(100.0)	
Total number of children wanted				14.1677
1-4	82 (32.4)	171 (67.6)	253	(<0.001) *
5+	244 (38.4)	392 (61.6)	(100.0)	
			636	
			(100.0)	
Gravida (number of pregnancies)				15.4730
1-4	236 (33.6)	467 (66.4)	703	(<0.001) *
5-9	98 (48.8)	103 (51.2)	(100.0)	
			201	
			(100.0)	
Age of current pregnancy(n=511)				14.0982
1 month	3 (37.5)	5 (62.5)	8 (100.0)	(0.119)
2 months	17 (47.2)	19 (52.8)	36 (100.0)	
3 months	21 (38.9)	33 (61.1)	54 (100.0)	
4 months	37 (52.1)	34 (47.9)	71 (100.0)	
5 months	34 (40.0)	51 (60.0)	85 (100.0)	
6 months	18 (29.5)	43 (70.5)	61 (100.0)	
7 months	25 (30.5)	57 (69.5)	82 (100.0)	
8 months	22 (29.3)	53 (70.7)	75 (100.0)	
9 months	10 (32.3)	21 (67.7)	31 (100.0)	
Post Date	3 (37.5)	5 (62.5)	8 (100.0)	
Ever had a childbirth complication(baby) (n=904)				0.6474
Yes	6 (28.6)	15 (71.4)	21 (100.0)	(0.421)
No	328 (37.1)	555 (62.9)	883	
			(100.0)	
Start of sexual intercourse after delivery (n=904)				42.8910
1-3 weeks after delivery	3 (33.3)	6 (66.7)	9 (100.0)	(<0.001) *
1-3 months after delivery	192 (48.0)	208 (52.0)	400	
4-6 months after delivery	62 (27.2)	166 (72.8)	(100.0)	
7-12 months after delivery	69 (32.1)	146 (67.9)	228	
>12 months	8 (15.4)	44 (84.6)	(100.0)	
			215	
			(100.0)	
			52 (100.0)	
Ever been counselled on FP? (n=904)				0.0444
Yes	243 (37.2)	411 (62.8)	654	(0.833)
No	91 (36.4)	159 (63.6)	(100.0)	
			250	
			(100.0)	
Ever used FP methods? (n=904)				0.444
Yes	243 (37.2)	411 (62.8)	654	(0.833)
No	91 (36.4)	159 (63.6)	(100.0)	
			250	

			(100.0)	
Ever used a FP method after delivery?				10.4212
	225 (41.1)	322 (58.9)	547	(0.001) *
Yes	109 (30.5)	248 (69.5)	(100.0)	
No			357	
			(100.0)	
Breastfed previous child? (n=904)				5.8653
Yes	319 (36.3)	560 (63.7)	879	(0.015) *
No	15 (60.0)	10 (40.0)	(100.0)	
			25 (100.0)	
Ever practiced exclusive breastfeeding? (n=904)				31.0284
Yes	230 (32.3)	482 (67.7)	712	(<0.001) *
No	104 (54.2)	88 (45.8)	(100.0)	
			192	
			(100.0)	
Duration of breastfeeding (n=712)				45.3461
less than 6 months	3 (7.3)	38 (92.7)	41 (100.0)	(<0.001) *
6 months to one year	135 (45.5)	162 (54.5)	297	
Above one year	92 (24.6)	282 (75.4)	(100.0)	
			374	
			(100.0)	
Ever experienced infertility issues? (n=904)				10.8256
Yes	3 (9.4)	29 (90.6)	32 (100.0)	(0.001) *
No	331 (38.0)	541 (62.0)	872	
			(100.0)	
Frequency of couples' sexual intercourse (n=904)				26.5771
Once a week	56 (29.2)	136 (70.8)	192	(<0.001) *
More than once in a week	163 (42.9)	217 (57.1)	(100.0)	
Once a month	15 (20.5)	58 (79.5)	380	
More than once in a month	93 (41.5)	131 (58.5)	(100.0)	
Once a year	6 (26.1)	17 (73.9)	73 (100.0)	
More than once in a year	1 (8.3)	11 (91.7)	224	
			(100.0)	
			23 (100.0)	
			12 (100.0)	

3.3 Logistic regression analysis of relationship between independent variables and BPI

Table 3 beneath shows the relationship between the independent variables and interbirth spacing. The binary logistic regression (BLR) analysis was carried out. In the BLR analysis, the factors that were significantly associated with interbirth spacing and those from the literature were sent to the multivariable logistic regression level for further analysis. After adjusting for confounders, women whose partners had higher educational attainment were twice as likely to space their births with no education as the reference (AOR 2.07; 95%CI1.09-3.96; p=0.027). Also,

women whose partners had secondary education were 1.87 times more likely to have longer birth intervals with reference to women whose partners had no education (AOR 1.87; 95%CI 1.1-3.17; p=0.021). We found that women who went through CS delivery were 3.28 times more likely to space their children compared to those who had SVD (AOR 3.28; 95% CI 1.02-10.62; p=0.047). It was further observed that multiparous women (five or more children) were 0.67 times more likely to have a longer BPI compared to women with less than five children (AOR 0.67; 95%CI 0.46-0.98; p=0.040). Finally, after the adjustment, it was found that the frequency of the couple's sexual intercourse was linked with BPI. The more frequent the sexual contacts, the shorter the interbirth interval.

Table 3. Logistic regression analysis of the relationship between independent variables and BPI

Independent variables	Interbirth interval between index child and subsequent pregnancy 1-23 months, n (%)		p-value	24-≥59 months, n (%)		
	Unadjusted Odds Ratio (95%CI)			Adjusted Odds Ratio (95% CI)	p-value	
Location/municipal						
BaM	Reference			Reference		
BoM	0.39	(0.2472-	(<0.001)	1.09	(0.267-0.4.474)	0.902
KNM	0.6068)	*		1.57	(0.310-7.947)	0.586
	0.55	(0.3472-	0.009			
		0.8578)				
Educational level of woman's partner						
No education	Reference			Reference		
Primary	0.85	(0.573-	0.414	1.39	(0.858-2.242)	0.182
JHS	1.258)		0.080	0.99	(0.619-1.598)	0.983
Secondary	0.71	(0.481-	0.698	1.87	(1.099-3.168)	0.021*
Higher	1.042)		0.100	2.07	(1.085-3.959)	0.027*
	1.09	(0.715-				
		1.651)				
	1.50	(0.925-				
		2.435)				
Religious affiliation						
Christian	Reference			Reference		
Islam	1.85	(1.26-2.704)	0.002*	1.15	(0.547-2.424)	0.710
Traditionalist	0.41	(0.156-	0.067	0.69	(0.242-1.978)	0.492
No religion	1.064)		0.432	1.03	(0.210-5.012)	0.973
	1.71	(0.449-				
		6.494)				
Ethnic background						
Frafra	Reference			Reference		
Kassena	1.20	(0.846-	0.305	0.83	(0.245-2.842)	0.771
Nankana	1.709)		0.131	0.53	(0.149-1.891)	0.329

Kusaasi	1.35	(0.915-	< 0.001 *	1.51 (0.299-7.609)	0.618
Builsa	1.986)		0.005 *	3.38 (0.559-20.446)	0.185
Moshie	3.24	(1.785-	0.014 *	2.97 (0.456-19.426)	0.255
Bisa	5.898)		0.392	0.66 (0.075-5.746)	0.704
Akan	5.74	(1.694-	0.859	0.59 (0.120-2.945)	0.523
Others	19.453)		0.285	0.27 (0.060-1.205)	0.086
	3.93	(1.324-			
	11.677)				
	1.68	(0.510-			
	5.568)				
	1.12	(0.312-			
	4.046)				
	0.62	(0.263-			
	1.480)				
Distance to Health Facility (HF)					
0–5km.	Reference			Reference	
6–10km	1.62	(1.201-	0.002 *	0.86 (0.506-1.446)	0.559
11–14km	2.188)		0.001 *	0.97 (0.467-2.016)	0.936
	2.05	(1.328-			
	3.150)				
Travel time to HF					
1–5min	Reference			Reference	
6–15min	0.94	(0.469-	0.850	0.75 (0.339-1.650)	0.472
16–29min	1.865)		0.442	1.03 (0.450-2.367)	0.941
30min-1hrs	1.32	(0.648-	0.051*	1.35 (0.527-3.478)	0.529
1hr–2hrs	2.706)		0.206	1.21 (0.310-4.762)	0.781
2hrs and above	2.10	(0.996-		1 (empty)	
	4.409)				
	1.96	(0.0.690-			
	5.590)				
	1 (empty)				
Means of transport to HF					
	Reference			Reference	
Bicycle	0.72	(0.258-	0.536	0.53 (0.151-1.833)	0.314
Car	2.024)		0.032 *	0.60 (0.328-1.092)	0.094
Foot	0.61	(0.384-	0.370	0.66 (0.358-1.234)	0.196
Motor bike	0.959)		0.340	0.88 (0.430-1.703)	0.724
Tricycle	0.81	(0.505-			
	1.289)				
	1.32	(0.746-			
	02.339)				
Reasons for preferred sex of child					
I have no preferred sex.	Reference			Reference	
Already had a girl (s).	1.41	(0.578-	0.449	0.76 (0.232-2.478)	0.647
Already had a boy (s).	3.448)		0.872	0.64 (0.194-2.129)	0.470
My husband's preferred choice	1.08	(0.426-	0.040 *	0.45 (0.194-1.029)	0.058
	2.739)		0.431	0.86 (0.423-1.703)	0.662
	0.49	(0.248-		1 (empty)	

Child was my preferred sex.	0.0.968) 0.83 (0.518-1.325) 1 (empty)			
Mode of delivery of last child	Reference		Reference	
SVD	3.76 (1.670-8.475)	0.001*	3.28 (1.015-10.616)	0.047*
CS	1 (empty)	0.357	1	
Vacuum extraction	1.57 (1.387-1.834)		1.18 (0.327-4.272)	0.798
Forceps delivery				
Parity	Reference		Reference	
1-3	0.67 (0.472-0.948)	0.024*	0.66 (0.433-1.009)	0.055
4-7				
Total number of children wanted	Reference		Reference	
1-4	0.57 (0.421-0.763)	<0.001*	0.67 (0.462-0.982)	0.040*
5+				
Start of sexual intercourse after delivery	Reference		Reference	
1-3 weeks after delivery	0.54 (0.134-0.2.196)	0.391	0.43 (0.077-2.385)	0.334
1-3 months after delivery	0.2.196)	0.686	1.02 (0.181-5.782)	0.979
4-6 months after delivery	1.34 (0.325-5.518)	0.938	1.85 (0.316-10.868)	0.494
7-12 months after delivery	5.518)	0.206	2.25 (0.330-15.375)	0.407
>12 months	1.06 (0.257-4.356) 2.75 (0.568-0.13.317)			
Frequency of couples' sexual intercourse	Reference		Reference	
Once a week	0.55 (0.378-0.0.795)	0.002*	0.67 (0.422-1.091)	0.109
More than once in a week	0.0.795)	0.159	0.96 (0.438-2.095)	0.914
Once a month	1.59 (0.833-0.3.042)	0.009*	0.35 (0.186-0.641)	0.001*
More than once in a month	0.3.042)	0.758	0.61 (0.191-1.915)	0.393
Once a year	0.58 (0.385-0.873)	0.153	2.59 (0.270-24.885)	0.409
More than once in a year	0.873)			
	1.17 (0.437-0.3.113) 4.53 (0.571-035.918)			

Figure 1: BPI between index child and subsequent pregnancy and maternal complications

As depicted in **Figure 1**, approximately, 9.5% (86/904) of the women reported experiencing complications during childbirth with retained product

of conception (41.9%), pregnancy induced hypertension (27.9%) and bleeding after delivery (11.6%) being the most commonly reported complications.

Figure 2: BPI between index child and subsequent pregnancy and perinatal complications

In terms of perinatal birth complications, 2.3% (21/904) of the respondents reported ever experiencing a complication including, neonatal sepsis (38.1%) and neonatal jaundice (23.8%); the rest are represented on the bar graph in figure 2 below.

Figure 1. Maternal birth complications during childbirth





Figure 2. Perinatal birth complications

4.0 Discussion

This study investigated the association between SBPI and perinatal and maternal outcomes among 904 women in three municipalities in the UER of Ghana. The prevalence of the SBPI (<24 months) was 36.9% among the respondents in this study. This proportion is quite lower than a study that was conducted in Ghana (Alhassan et al., 2022), which showed a SBPI of 49.7%. The disparities could be due to variations in sample size between the two studies. Again, this finding is in line with recent studies that reveal similar but higher SBPI prevalence in Sudan (60.6%) (Damtie et al., 2021), Ethiopia (58.74%) (Belachew et al., 2023), and Uganda (52.4%) (Aleni et al., 2020). Much lower prevalence was found in studies conducted in Iran (28.8%) and rural Bangladesh (24.6%).

The results also revealed that BPI was significantly associated with some adverse maternal and perinatal consequences. However, there was not much difference in terms of complications experienced between women who had a SBPI (<24 months) and the optimal interval (24–59 months). Eight percent (8.1%) of the women who had a SBPI (<24 months) reportedly experienced birth complications, while 10.4% of them within the interval of 24–59 months reported birth complications. The major birth complications encountered by women during childbirth were retained products of

conception (41.9%), pregnancy-induced hypertension (27.9%), postpartum hemorrhage (11.6%), and obstructed labor (10.5%). Similarly, sepsis (38.1%), neonatal jaundice (23.8%), low birth weight (19%), preterm birth (14.3%), and early neonatal death were the reported perinatal complications in this study. Previous studies have shown resemblance maternal and perinatal complications. For instance, a systematic review study (Hutcheon et al., 2019) in 2019 indicated that inadequately spaced births result in birth complications. A recent meta-analysis research revealed that women with SBPI had nearly 1.5 times the odds of developing either a preterm or low-birthweight infant (David et al., 2017). A United States study (Byamukama et al., 2022) shown that SBPI is associated with premature rupture of membranes. Another systematic review study in China (Xu et al., 2022) found that short birth spacing was associated with an increased risk of uterine rupture, placenta previa, and placental abruption. A prior study (Rao et al., 2022) conducted in 2022 indicated low birth weight (OR 3.5, 95% CI:1.2–10.3, $P < 0.05$), preterm birth (OR 5.5, 95% CI:1.5–21.3, $P < 0.05$), and postpartum hemorrhage (OR 19.6, 95% CI:4.4–90.9, $P < 0.05$), were significantly higher in respondents with SBPI than in those with an OBPI (24–59 months).

The educational level of women partner (AOR 2.07; 95% CI 1.09–3.96; $p=0.027$), mode of delivery of the last child (AOR 3.28; 95% CI 1.02–10.62; $p=0.047$), parity of five or more children (AOR 0.67; 95% CI 0.46–0.98; $p=0.040$) and frequency of sexual intimacy were strongly associated with BPI after adjusting for confounders. Women whose partners had secondary education were 1.87 times more likely to space their births compared to those with partners without education. This result confirms with studies that were conducted in Ethiopia (Hailu & Gulte, 2016), and Korea. However, unlike our study, these two studies and many more found an association with women's education with BPI instead. A survey among college students postulated that couples with lower or no education had a lesser chance of spacing their births than educated ones (Shen, 2019). This could be due to the fact that educated females are more likely to utilize birth control methods to lengthen their birth intervals and have available health information. Similarly, a recent study opined that mothers with low education levels are at increased risk of SBPI (Pimentel et al., 2020).

We found that women who went through CS during delivery were 3.28 times more likely to space their children compared to those who had SVD. This finding is in conformity with a study (Byamukama et al., 2022), which stated that long birth intervals were significantly associated with CS (0.69; 0.56–0.82), which have a greater risk of premature rupture of membranes (PROM) (1.57; 1.20–2.06) and low birth weight (1.46; 1.03–2.06). This revelation could be linked to the fact that when women undergo

an operation during childbirth, they need more time recover before the initiation the next pregnancy, especially in the case of birth complications(Huber et al., 2020).

In terms of parity, this study found a significant positive impact on the risk of having a subsequent birth. Study results are associated with this present study, such that a study in Manipur and Babol, Northern Iran, indicated that birth interval increases with increasing parity (Nausheen et al., 2021). This seeks to explain that women with fewer children tend to have short intervals while those with more children have longer intervals. The association between the number of children and women whose child dies immediately after birth has shorter intervals. For instance, a study in Poland by Merklinger-Gruchala *et al.* revealed that a woman who gives birth to a preterm baby that subsequently dies will again try to become pregnant more quickly. Therefore, her BPI becomes very short.

It was further discovered in this study that couples who had more frequent sexual contacts had a high probability of having a shorter interbirth spacing compared to couples who had less frequent intimacy. It is assumed that when couples engage in frequent sexual intimacy, could easily result in pregnancy, hence a shorter BPI. This is valid for couples who live together in the same household. The majority of respondents in this study lives in the same households with their partners. This further explains that couples using modern contraceptives have a higher percentage of widening their birth interval than those who are not practicing any form of FP. This finding is in line with a study(Yussuf et al., 2020;Zimmerman et al., 2019) in Tanzania and Ethiopia respectively.

This study result should be interpreted carefully since it is still unclear whether the link between SBPI and adverse health outcomes is causal or confounded by differences in the health characteristics. Further research is therefore needed to determine if the association between SBPI and adverse health outcomes is due to a direct cause-and-effect relationship or if it is influenced by other factors such as lifestyle choices or access to healthcare. Understanding these potential confounding variables will provide a more comprehensive understanding of the relationship between SBPI and adverse health outcomes.

4.1 *Strengths and Limitations of the Study*

The study has some limitations. The sample sizes for the three study sites were unequal due to a dropout rate at the BaM study site. This was due to an outbreak of conflict during the period of data collection in the area. Again, the study did not conform strictly with the inclusion criteria (women with two successive live births prior to data collection). A few uniparous women were included in the study.

4.2 **Conclusion and Global Health Implications**

The study evidence prongs to critical periods at which women should avoid getting pregnant to preempt poor prenatal and maternal outcomes associated with inappropriate birth intervals. The study recommends health professionals, especially those in direct contact with family, maternal, reproductive, and child health, to identify the challenges facing contraceptive uptake particularly among multiparous women and intensify birth spacing campaigns to improve optimal birth spacing and to support women's decision-making.

Maternal and newborn mortalities in Ghana and SSA remain radically high, while contraceptive uptake has dwindled. Hence, this study findings provides empirical evidence for policy interventions to foster optimal birth spacing to ensure a reduction in high fertility and maternal and child mortality.

Conflicts of Interest: The authors declare no conflict of interest in the conduct of this study.

Financial Disclosure: There is nothing to declare.

Funding/Support: There was no funding for this study.

Ethics Approval: Ethical clearance was sought from the Navrongo Health Research Centre Institutional Review Board of Ghana and the Ghana Health Service through the Upper East Regional Health Directorate. Written permission to carry out the study was sought from the municipal health authorities.

References:

1. Ajayi, A. I., & Somefun, O. D. (2020). Patterns and determinants of short and long birth intervals among women in selected sub-Saharan African countries. *Medicine*, 99(19), e20118. <https://doi.org/10.1097/MD.00000000000020118>
2. Alhassan, A. R., Anyinzaam-Adolipore, J. N., & Abdulai, K. (2022). Short birth interval in Ghana: Maternal socioeconomic predictors and child survival. *Population Medicine*, 4(January), 1–8. <https://doi.org/10.18332/POP/145914>
3. Ayane, G. B., Desta, K. W., Demissie, B. W., Assefa, N. A., & Woldemariam, E. B. (2019). Suboptimal child spacing practice and its associated factors among women of child bearing age in Serbo town, JIMMA zone, Southwest Ethiopia. *Contraception and Reproductive Medicine*, 4(1), 1–8. <https://doi.org/10.1186/s40834->

019-0085-1

4. Bauserman, M., Nowak, K., Nolen, T. L., Patterson, J., Lokangaka, A., Tshefu, A., Patel, A. B., Hibberd, P. L., Garces, A. L., Figueroa, L., Krebs, N. F., Esamai, F., Liechty, E. A., Carlo, W. A., Chomba, E., Mwenechanya, M., Goudar, S. S., Ramadurg, U., Derman, R. J., ... Bose, C. (2020). The relationship between birth intervals and adverse maternal and neonatal outcomes in six low and lower-middle income countries. *Reproductive Health*, *17*(Suppl 2), 1–10. <https://doi.org/10.1186/s12978-020-01008-4>
5. Belachew, T. B., Asmamaw, D. B., & Negash, W. D. (2023). Short birth interval and its predictors among reproductive age women in high fertility countries in sub-Saharan Africa: a multilevel analysis of recent Demographic and Health Surveys. *BMC Pregnancy and Childbirth*, *23*(1), 1–9. <https://doi.org/10.1186/s12884-023-05403-0>
6. Byamukama, O., Migisha, R., Kalyebara, P. K., Tibajjuka, L., Lugobe, H. M., Ngonzi, J., Ahabwe, O. M., Garcia, K. R. M., Mugenyi, G. R., Boatun, A. A., Muhumuza, J., Ssalongo, W. G. M., Kayondo, M., & Kanyesigye, H. (2022). Short interbirth interval and associated factors among women with antecedent cesarean deliveries at a tertiary hospital, Southwestern Uganda. *BMC Pregnancy and Childbirth*, *22*(1), 1–8. <https://doi.org/10.1186/s12884-022-04611-4>
7. Conde-Agudelo, A., Rosas-Bermudez, A., & Norton, M. H. (2016). Birth spacing and risk of autism and other neurodevelopmental disabilities: A systematic review. *Pediatrics*, *137*(5). <https://doi.org/10.1542/peds.2015-3482>
8. Damtie, Y., Kefale, B., Yalew, M., Arefaynie, M., & Adane, B. (2021). Short birth spacing and its association with maternal educational status, contraceptive use, and duration of breastfeeding in Ethiopia. A systematic review and metaanalysis. *PLoS ONE*, *16*(2 February), 1–15. <https://doi.org/10.1371/journal.pone.0246348>
9. David, H. C., Amani, M. N.-J., & Nancy, E. A. (2017). 乳鼠心肌提取 HHS Public Access. *Physiology & Behavior*, *176*(3), 139–148. <https://doi.org/10.1038/s41372-019-0402-1>.Short
10. Dehesh, T., Salarpour, E., Malekmohammadi, N., & Kermani, S. A. (2020). Associated factors of pregnancy spacing among women of reproductive age Group in South of Iran: Cross-sectional study. *BMC Pregnancy and Childbirth*, *20*(1), 1–7. <https://doi.org/10.1186/s12884-020-03250-x>
11. Guure, C., Maya, E. T., Dery, S., Da-Costa Vrom, B., Alotaibi, R. M., Rezk, H. R., & Yawson, A. (2019). Factors influencing unmet need for family planning among Ghanaian married/union women: A multinomial mixed effects logistic regression modelling approach.

- Archives of Public Health*, 77(1), 1–12.
<https://doi.org/10.1186/s13690-019-0340-6>
12. Hailu, D., & Gulte, T. (2016). Determinants of Short Interbirth Interval among Reproductive Age Mothers in Arba Minch District, Ethiopia. *International Journal of Reproductive Medicine*, 2016, 1–17. <https://doi.org/10.1155/2016/6072437>
 13. Huber, L. R. B., Smith, K., Sha, W., & Vick, T. (2020). *Interbirth Interval and Pregnancy Complications and Outcomes: Findings from the Pregnancy Risk Assessment Monitoring System*. 1–10. <https://doi.org/10.1111/jmwh.12745>
 14. Hutcheon, J. A., Nelson, H. D., Stidd, R., Moskosky, S., & Ahrens, K. A. (2019). Short interpregnancy intervals and adverse maternal outcomes in high-resource settings: An updated systematic review. *Paediatric and Perinatal Epidemiology*, 33(1), O48–O59. <https://doi.org/10.1111/ppe.12518>
 15. Mehra, R., Keene, D. E., Kershaw, T. S., Ickovics, J. R., & Warren, J. L. (2019). Racial and ethnic disparities in adverse birth outcomes: Differences by racial residential segregation. *SSM - Population Health*, 8(January), 2–11. <https://doi.org/10.1016/j.ssmph.2019.100417>
 16. Merdad, L., & Ali, M. M. (2018). *Timing of maternal death : Levels , trends , and ecological correlates using sibling data from 34 sub-Saharan African countries*. 5, 1–13.
 17. Merklinger-Gruchala, A., Jasienska, G., & Kapiszewska, M. (2015). Short interpregnancy interval and low birth weight: A role of parity. *American Journal of Human Biology*, 27(5), 660–666. <https://doi.org/10.1002/ajhb.22708>
 18. Molitoris, J. (2018). Heterogeneous Effects of Birth Spacing on Neonatal Mortality Risks in Bangladesh. *Studies in Family Planning*, 49(1), 3–21. <https://doi.org/10.1111/sifp.12048>
 19. Nausheen, S., Bhura, M., Hackett, K., Hussain, I., Shaikh, Z., Rizvi, A., Ansari, U., Canning, D., Shah, I., & Soofi, S. (2021). Determinants of short birth intervals among married women: A cross-sectional study in Karachi, Pakistan. *BMJ Open*, 11(4), 1–10. <https://doi.org/10.1136/bmjopen-2020-043786>
 20. Pimentel, J., Ansari, U., Omer, K., Gidado, Y., Baba, M. C., Andersson, N., & Cockcroft, A. (2020). Factors associated with short birth interval in low- And middle-income countries: A systematic review. *BMC Pregnancy and Childbirth*, 20(1), 1–17. <https://doi.org/10.1186/s12884-020-2852-z>
 21. Rao, J., Fan, D., Ma, H., Lin, D., Zhang, H., Zhou, Z., Li, P., Chen, G., Lu, D., Liu, Y., Wu, Z., He, J., Liu, X., Peng, B. J., Guo, X., &

- Liu, Z. (2022). Is there an optimal inter-delivery interval in women who underwent trial of labor after cesarean delivery (TOLAC)? *Reproductive Health*, *19*(1), 1–7. <https://doi.org/10.1186/s12978-021-01319-0>
22. Schummers, L., Hutcheon, J. A., Hernandez-Diaz, S., Williams, P. L., Hacker, M. R., Vanderweele, T. J., & Norman, W. V. (2018). Association of Short Interpregnancy Interval with Pregnancy Outcomes According to Maternal Age. *JAMA Internal Medicine*, *178*(12), 1661–1670. <https://doi.org/10.1001/jamainternmed.2018.4696>
23. Shen, D. (2019). *Munich Personal RePEc Archive Better Educated , Fewer Divorces : The Impact of College Education Quality on Marriage Outcomes Better Educated , Fewer Divorces : The Impact of College Education Quality on Marriage Outcomes*. 94198.
24. Stover, J., & Winfrey, W. (2017). The effects of family planning and other factors on fertility, abortion, miscarriage, and stillbirths in the Spectrum model. *BMC Public Health*, *17*(Suppl 4). <https://doi.org/10.1186/s12889-017-4740-7>
25. Survey, M. H., & Indicators, K. (2012). Key Indicators. *Timing Solutions for Swing Traders*, 91–111. <https://doi.org/10.1002/9781119199045.ch6>
26. Tesema, G. A., Worku, M. G., & Teshale, A. B. (2021). Duration of birth interval and its predictors among reproductive-age women in Ethiopia: Gompertz gamma shared frailty modeling. *PLoS ONE*, *16*(2 February 2021), 1–13. <https://doi.org/10.1371/journal.pone.0247091>
27. Tessema, Z. T., Yazachew, L., Tesema, G. A., & Teshale, A. B. (2020). Determinants of postnatal care utilization in sub-Saharan Africa: a meta and multilevel analysis of data from 36 sub-Saharan countries. *Italian Journal of Pediatrics*, *46*(1), 1–11. <https://doi.org/10.1186/s13052-020-00944-y>
28. Xu, T., Miao, H., Chen, Y., Luo, L., Guo, P., & Zhu, Y. (2022). Association of Interpregnancy Interval With Adverse Birth Outcomes. *JAMA Network Open*, *5*(6), E2216658. <https://doi.org/10.1001/jamanetworkopen.2022.16658>
29. Yussuf, M. H., Elewonibi, B. R., Rwabilimbo, M. M., Mboya, I. B., & Mahande1, M. J. (2020). Trends and predictors of changes in modern contraceptive use among women aged 15–49 years in Tanzania from 2004–2016: Evidence from Tanzania Demographic and Health Surveys. *PLoS ONE*, *15*(6 June), 1–14. <https://doi.org/10.1371/journal.pone.0234980>
30. Zimmerman, L. A., Yi, Y., Yihdego, M., Abrha, S., Shiferaw, S., Seme, A., & Ahmed, S. (2019). Effect of integrating maternal health

services and family planning services on postpartum family planning behavior in Ethiopia: Results from a longitudinal survey. *BMC Public Health*, 19(1), 1–9. <https://doi.org/10.1186/s12889-019-7703-3>