

Selection and Phenotypic Stability of M4 Mutants of Pearl Millet (*Pennisetum glaucum* (L.) R. Br.) Derived from Gamma Rays Induced Mutagenesis

Mouhamadou Mounkaila Boureima

Sani Daouda Ousmane

Abdoul-Bachir Issa Yacouba

Abdoul Razak Sani Daoud

Nassirou Mahamadou Adamou

Mamane Nassourou Lawali

Illiassa Soumaila Sounakoye

Université Abdou Moumouni, Institut des Radio Isotopes, Niamey, Niger

[Doi: 10.19044/esipreprint.9.2024.p124](https://doi.org/10.19044/esipreprint.9.2024.p124)

Approved: 05 September 2024

Posted: 08 September 2024

Copyright 2024 Author(s)

Under Creative Commons CC-BY 4.0

OPEN ACCESS

Cite As:

Mounkaila Boureima M., Daouda Ousmane S., Issa Yacouba A.B., Sani Daoud A.R., Mahamadou Adamou N., Lawali M.N. & Soumaila Sounakoye I. (2024). *Selection and Phenotypic Stability of M4 Mutants of Pearl Millet (Pennisetum glaucum (L.) R. Br.) Derived from Gamma Rays Induced Mutagenesis*. ESI Preprints.

<https://doi.org/10.19044/esipreprint.9.2024.p124>

Abstract

Gamma irradiation mutagenesis is an approach that offers a wide range of possibilities for varietal selection. It consists of simultaneously inducing multiple mutations to modify several plant traits. The aim of the present study was to select seed lots for four mutants in the M4 generation. An experiment was carried out in an experimental plot to purify drought-tolerant M4 genotypes based on morphological characteristics. Potential mutants MI 02/82, MI 13/63, MI 12/72 and MI 10/54 were tested in a randomized complete block design. The parameters measured are the number of tillers, stem height, number of internodes, number of ears, ear length, stem and spike diameters and cycle duration. The results show on morphological parameters that MI 13/63 and MI 10/54 genotypes as an homogeneous population from M4 onwards. MI 12/72 can be classified into two different subgroups according to ear length. MI 02/82, on the other hand, shows a high degree of variability at M4. These results will contribute to the selection of

new varieties adapted to the requirements of rural producers to improve pearl millet productivity in Niger.

Keywords: Pearl millet, Gamma rays, M4 mutants, Crop breeding, Morphological diversity

Introduction

Millet [*Pennisetum glaucum* (L.) R. Br.] is the sixth most important cereal in the world, mainly cultivated on thirty million hectares in the arid and semi-arid zones of Asia and Africa (Satyavathi et al., 2021). Millet cultivation is facing the effects of climate change, which induce losses of up to 20% in grain and 5% on biomass (Benoit & Mandéla, 2015). The vegetative, reproductive and physiological characteristics of millet give it tolerance to low soil fertility, high soil pH, drought, high temperatures and salinity (Varshney et al., 2017). Selection by mutagenesis is a flexible and practical approach, applicable to any crop with precise objectives and appropriate selection methods (Oladosu et al., 2016). It has become an interesting approach to plant improvement and its advantage lies in its ability to improve one or more specific traits of the preferred variety (Hase et al., 2020). Gamma rays have the advantage of being highly effective in generating a high frequency of mutations that can lead to loss of gene function (Kazama et al., 2017).

This is why gamma rays have become the most frequently used mutagenic agent in plant breeding (Beyaz & Yildiz, 2017; Hase et al., 2020). However, in millet breeding, few studies have been carried out on the rate and effectiveness of physical mutagens on the characteristics of inbred millet lines due to natural genetic variation (Serba et al., 2017). The effects of gamma rays mutagenesis on the morphological parameters of millet are multiple and manifest themselves on the height and diameter of stems, leaf shrinkage, seed size, earliness or lengthening of the cycle and the length of ears (Ambli & Mullainathan, 2015). These mutations are characterized by a high heritability of characters in M2 (Abdalla et al., 2016) and on the number of nodes, stem and ear diameter in the M3 generation (Ambli, 2018). However, selection in the M3 generation of millet mutants is less effective because of the high genetic variability. (Maryono et al., 2020).

As part of the genetic improvement program for water stress tolerance, the HKP (Haini Kirey Precoce) millet variety was irradiated with 250 Gray gamma rays. In morphological terms, HKP has an average tillering capacity, with stems 190 to 200 cm long, and ears of intermediate length. It has the particularity of being an earlier local variety, with a sowing-maturing cycle of 75 to 90 days (MA/Niger, 2021). The aim of this study is to

investigate the morphological variability of M4 genotypes for the effective selection of four water-stress tolerant millet mutants.

Methods

The material consists of populations of four drought-tolerant M4 genotypes MI 02/82, MI 13/63, MI 12/72 and MI 10/54 of millet derived from a mutagenesis process using gamma irradiation at *Laboratoire de Biotechnologie et Amélioration des plantes (LABAP)*. The device is a randomized complete block (Figure 1). All measures were taken to ensure controlled pollination. The following parameters were measured: heading date (HD), number of tillers (NT) per plant, stem diameter (SD) and ear diameter (ED), plant height (PLH) and ear length (EL), number of ears (NE) and number of nodes (NN). A hierarchical classification was carried out to generate dendrograms in the populations of each of the four genotypes, using R software. Correlations between measured parameters were highlighted by bottom-up classification. Branches at the precise height of 2 dendrogram units were taken as the discrimination threshold. The averages of individuals from the different class, were analyzed by ANOVA (at the 5% threshold) using GenStat software, to highlight their homogeneity.




Figure 1 : Schematic diagram of the experimental set-up. Each block (A, B, C and D) is dedicated to a single genotype family

Results

Analysis of genotype diversity based on dendrograms reveals a higher height for MI 02/82 and MI 13/63 (10 and 7 units) than for MI 12/72 and MI 10/54 (6 and 5 units) (Figure 2). At the precise height of 2 units on the dendrogram scale in Figure 2a, MI 02/82 genotypes hold more branches (13 branches) compared with only 6, 5 and 4 branches for MI 13/63, MI 12/72 and MI 10/54 respectively (Figure 2b, c and d). Through the subfamilies, MI 13/63 and MI 10/54 genotypes have 10.7 and 8.5 nodes respectively for stem heights of 182.8 and 157 cm. Whereas MI 12/72 has an average of 8.8 nodes for a stem height of 150 cm.

The results of the analysis of variance allow us to transform the branches of the MI 13/63, MI 12/72 and MI 10/54 genotypes into different classes, based on the high similarity between individuals. This allows us to distinguish 13 class for MI 02/82, 6 class for MI 13/63, 5 and 4 class, respectively for MI 12/72 and MI 10/54.

These results indicate MI 10/54 genotypes, that are statically identical for all the parameters studied. The MI 13/63 and MI 12/72 genotypes were different in number of tillers (Figure 3) and ear length (Figure 4), respectively. For MI 13/63 genotype, the difference between subgroups was in the number of tillers, which varied from 9.5 to 19.5 tillers (Figure 3). Whereas in MI 12/72, ear length varies with an average of 36 to 52 cm between classes (Figure 4). Contrary to that MI 02/82 classes are statistically different on millet varietal characteristics (Figure 5A-D).




Figure 2 : Hierarchical classification of genotypes. *The letters a, b, c and d, designate the MI 02/82, MI 13/63, MI 12/72 and MI 10/54 dendrograms respectively.*

Table 4 : Classification of MI 10/54 genotypes into subfamilies

Parameters	Class					
	D1	D2	D3	D4	Means	P Value*
HD	65.5 ±2.1	56.5 ±2.1	52.0 ±2.8	61.5 ±9.2	59.0	0.283
NN	9.5 ±0.7	7.0 ±0.0	9.0 ±0.0	8.5 ±0.7	8.5	0.072
NT	18.5 ±2.1	15.0 ±0.0	13.0 ±4.2	13.0 ±0.0	14.9	0.179
NE	9.5 ±4.9	5.0 ±2.8	8.0 ±1.4	6.5 ±0.7	7.2	0.558
SD	1.3 ±0.3	0.8 ±0.0	1.1 ±0.1	1.5 ±0.0	1.2	0.068
PLH	168.5 ±16.3	140.0 ±28.3	158.5 ±2.1	161.0 ±5.7	157.0	0.419
EL	36.5 ±7.8	36.5 ±3.5	385 ±0.7	57.0 ±9.9	42.1	0.052
ED	1.5 ±0.1	1.7 ±0.0	1.8 ±0.2	2.0 ±0.2	1.8	0.187

HD : Heading date, *NN* : Number of nodes, *NT* : Number of tillers, *NE* : Number of ears, *SD* : Stem diameter, *PLH* : Plant height, *EL* : Ear length, *ED* : Ear diameter. *If P value is less than 0.05, the difference between the means is significant






Figure 3 : MI 13/63 number of tiller (NT) classification ($p < 0,001$). The letters *a* and *b* indicate significantly different classes for the number of tillers Parameter.

Figure 4 : MI 12/72 ear length classification (EL) ($p = 0,001$). The letters *a* and *b* indicate significantly different classes for the ear length parameter

Figure 5 : Variability in the morphological characteristics of MI 02/82

Letter “A” indicates the heading date (HD), letter “B” indicates ear length (EL), letter “C” represents ear diameter (ED) and letter “D” illustrates stem diameter (SD). The letters a, b, c, d, e and f show the different classes in the analysis of variance at the 5% threshold

Discussion

Mutagenesis by gamma irradiation induces genetic variability in millet, making it possible to obtain mutants with outstanding agronomic performance (Addai & Yahaya, 2018). Selection significantly reduces genotypic and phenotypic variance from the M4 generation of gamma irradiated sesame mutants (ARISTYA et al., 2017). In millet, the number of effective stems per plant, panicle length and panicle diameter have a high heritability and genetic advance (PA et al., 2023). In this study, analysis by hierarchical classification shows morphological diversity that varies according to genotyp. All four genotype families performed individually. MI 02/82 shows more variability at generation M4 than genotypes MI 13/63, MI 12/72 (Figure 2). MI 10/54 genotype represents the most homogeneous of all the parameters studied in M4 generation (Table 4). Morphological variability in gamma-irradiated genotypes varies with dose and variety (Saibari et al., 2023). Because irradiation mutation randomly affects morphological characteristics (Abdullah et al., 2018). All four genotype families performed individually. This confirms the existence of intraspecific variability in crop radio-sensitivity reported by Hazra et al., (2022).

The MI 13/63 genotype family could be considered a morphologically homogeneous group. However, the number of tillers contributes significantly to the agronomic traits of millet. (Naoura et al.,

2020), but has low heritability (Yahaya, 2015). In rice M4 mutant, irradiation did not affect the number of tillers (Bella, 2021). On contrary, the average length of the ears is a selection index for millet (Patil et al., 2018). This makes it possible to distinguish two subgroups of MI 12/72 (short ears and medium ears) with a significant difference in ear length. Based on the classification of Singh et al., (2016), the MI 13/63 and MI 10/54 genotypes have a low number of nodes and medium stem height (Tables S2 and 4). Whereas MI 12/72 has short stems. This indicates that the MI 13/63 and MI 10/54 genotypes are similar in stem height to HKP, while MI 12/72 is shorter. This reduction in stem height could be induced by gamma irradiation. Similar effects were observed on some thirty M4 rice mutants (Hanifah et al., 2020).

Conclusions

In conclusion, the morphological purification of genotypes obtained by irradiation in millet depends on other factors, such as the nature of the mutation type, in addition to the species. Genotypes MI 13/63 and MI 10/54 are homogeneous, MI 12/72 has an intermediate homogeneity and MI 02/82 shows variability in all the parameters studied. MI 12/72 and MI 02/82 genotypes must be produced in M5 generation for purification. The homogeneity test can be tested on the M4 generation of pearl millet for gamma-induced mutation improvement.

Acknowledgments: The authors would like to thank Djatta Moumouni and all the students for their enormous contribution to data collection.

Conflict of Interest: The authors reported no conflict of interest.

Data Availability: All data are included in the content of the paper.

Funding Statement: The authors did not obtain any funding for this research.

References:

1. Abdalla, M., Yassein, H.E., & Okasha, M. (2016). Mutagenic effect of dimethyl sulphate and gamma rays on pearl millet (*Pennisetum glaucum*, L.). *Al-Azhar Journal for Agricultural Researches*, 26, 293-307.
2. Abdullah, S., Kamaruddin, N. Y., & Harun, A. R. (2018). The Effect of Gamma Radiation on Plant Morphological Characteristics of *Zingiber officinale* Roscoe. *International Journal on Advanced*

- Science, Engineering and Information Technology*, 8(5), 2085-2091. <https://doi.org/10.18517/ija.seit.8.5.4641>.
3. Addai, I., & Yahaya, B. (2018). Response of pearl millet (*Pennisetum glaucum*.) to experimental mutagenesis in the Guinea Savannah agro-ecology of Ghana. *Ghana Journal of Development Studies*, 15, 22. <https://doi.org/10.4314/gjds.v15i1.2>.
 4. Ambli, K., & Mullainathan, L. (2015). Chlorophyll and morphological mutants of Pearl millet (*Pennisetum typhoides* (Burn.) stapf. Var. CO (cu)9. Euro. J. Exp. Bio., 2015, 5(3) :72-77.
 5. Ambli, L. (2018). Effect of mutagens on quantitative characters in M3 generation of pearl millet (*Pennisetum typhoides* (BURN.F) STAPF. AND C.E. HUBB.). *Journal of Phytology*, 01-05. <https://doi.org/10.25081/jp.2018.v10.3404>.
 6. ARISTYA, V. E., Taryono, & Rani, A. W. (2017). Evaluation of genetic parameters in M4 and M5 generations of sesame mutant lines. *SABRAO J. Breed. Genet.* 49 (2) 201-201 https://acadstaff.ugm.ac.id/karya_files/evaluation-of-genetic-parameters-in-m4-and-m5-generations-of-sesame-mutant-lines-28f179c7eedf73fbb790ac461804b086.
 7. Bella, R. A. (2021). Improvement of Rice Phenotype of M4 Sigupai Irradiated by Gamma Ray. *Journal of Agriculture and Veterinary Science (IOSR-JAVS)* 2319-2372. Volume 14, Issue 3 Ser. II PP 36-41. <https://doi.org/10.9790/2380-1403023641>.
 8. Benoit, S., & Mandéla, H. (2015). Atlas agroclimatique sur la variabilité et le changement climatique au Niger (p. 37). Agrhymet /CILLS. <https://duddal.org/files/original/863a2c907349898300cea1414798ffc0dc0d56fa.pdf>.
 9. Beyaz, R., & Yildiz, M. (2017). The Use of Gamma Irradiation in Plant Mutation Breeding. In S. Jurić (Éd.), *Plant Engineering. InTech*. <https://doi.org/10.5772/intechopen.69974>.
 10. Hanifah, W. N., Parjanto, P., Hartati, S., & Yunus, A. (2020). The performances of M4 generation of Mentik Susu rice mutants irradiated with gamma-ray. *Biodiversitas Journal of Biological Diversity*, 21(9), Article 9. <https://doi.org/10.13057/biodiv/d210915>.
 11. Hase, Y., Satoh, K., Seito, H., & Oono, Y. (2020). Genetic Consequences of Acute/Chronic Gamma and Carbon Ion Irradiation of *Arabidopsis thaliana*. *Frontiers in Plant Science, Front Plant Sci.* 25 :11:336. doi: 10.3389/fpls.2020.00336. eCollection 2020.
 12. Hazra, S., Gorai, S., Bhattacharya, S., Bose, S., Hazra, P., Chattopadhyay, A., & Maji, A. (2022). Radio-sensitivity of diverse tomato genotypes with respect to optimization of gamma irradiation

- dose. *Brazilian Journal of Botany*, 45(3), 917-927. <https://doi.org/10.1007/s40415-022-00823-2>.
13. Kazama, Y., Ishii, K., Hirano, T., Wakana, T., Yamada, M., Ohbu, S., & Abe, T. (2017). Different mutational function of low- and high-linear energy transfer heavy-ion irradiation demonstrated by whole-genome resequencing of Arabidopsis mutants. *The Plant Journal*, 92(6), 1020-1030. <https://doi.org/10.1111/tpj.13738>.
 14. MA/Niger. (2021). Catalogue National des Espèces et Variétés Végétales (2 ; p. 304). Ministère de l'Agriculture. https://reca-niger.org/IMG/pdf/catalogue_des_especes_et_varietes_niger.pdf.
 15. Maryono, M., Sihono, Indriatama, W. M., & Human, S. (2020). Performance and estimation genetic variability of M3 pearl millet (*Pennisetum glaucum*) populations. *IOP Conference Series: Earth and Environmental Science*, 484(1), 012021. <https://doi.org/10.1088/1755-1315/484/1/012021>.
 16. Naoura, G., Sawadogo, N., Djirabaye, N., & Hassane, M. (2020). Agronomic performance of improved pearl millet cultivars in southern Chad. *International Journal of Biological and Chemical Sciences*, 14, 2980-2991. <https://doi.org/10.4314/ijbcs.v14i9.2>.
 17. Oladosu, Y., Rafii, M. Y., Abdullah, N., Hussin, G., Ramli, A., Rahim, H. A., Miah, G., & Usman, M. (2016). Principle and application of plant mutagenesis in crop improvement: A review. *Biotechnology & Biotechnological Equipment*, 30(1), 1-16. <https://doi.org/10.1080/13102818.2015.1087333>.
 18. PA, G., HS, P., PR, P., & Donga, A. (2023). Study of genetic variability, heritability and genetic advance for yield and its component traits in pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *The Pharma Innovation*, 12, 4305-4308. <https://doi.org/10.22271/tpi.2023.v12.i3au.19425>.
 19. Patil, K. S., Gupta, S., Dangi, K., Shashibhushan, D., Balram, M., & Ramesh, T. (2018). Panicle Traits and Plant Height are Important Selection Indices to Enhance Productivity in Pearl Millet (*Pennisetum glaucum* L.R.Br.) Populations. *International Journal of Current Microbiology and Applied Sciences*, 7, 306-312. <https://doi.org/10.20546/ijcmas.2018.712.037>.
 20. Saibari, I., Barrijal, S., Mouhib, M., Belkadi, N., & Hamim, A. (2023). Gamma irradiation-induced genetic variability and its effects on the phenotypic and agronomic traits of groundnut (*Arachis hypogaea* L.). *Frontiers in Genetics*, 14. <https://www.frontiersin.org/articles/10.3389/fgen.2023.1124632>.
 21. Satyavathi, C. T., Ambawat, S., Khandelwal, V., & Srivastava, R. K. (2021). Pearl Millet : A Climate-Resilient Nutricereal for Mitigating

- Hidden Hunger and Provide Nutritional Security. *Frontiers in Plant Science*, 12. <https://www.frontiersin.org/articles/10.3389/fpls.2021.659938>.
22. Serba, D. D., Perumal, R., Tesso, T. T., & Min, D. (2017). Status of Global Pearl Millet Breeding Programs and the Way Forward. *Crop Science*, 57(6), 2891-2905. <https://doi.org/10.2135/cropsci2016.11.0936>.
 23. Singh, S., Y, P., H, P., D, V., & Yadav, N. (2016). Morphological characterization of pearl millet hybrids [*Pennisetum glaucum* (L.) R. Br.] and their parents. *African Journal of Agricultural Research*, 11, 371-378. <https://doi.org/10.5897/AJAR2015.10333>.
 24. Varshney, R. K., Shi, C., Thudi, M., Mariac, C., Wallace, J., Qi, P., Zhang, H., Zhao, Y., Wang, X., Rathore, A., Srivastava, R. K., Chitikineni, A., Fan, G., Bajaj, P., Punnuri, S., Gupta, S. K., Wang, H., Jiang, Y., Couderc, M., Xu, X. (2017). Pearl millet genome sequence provides a resource to improve agronomic traits in arid environments. *Nature Biotechnology*, 35(10), 969-976. <https://doi.org/10.1038/nbt.3943>.
 25. Yahaya, Y. (2015). Correlation and Heritability in Pearl Millet (*Pennisetum glaucum* (L.) R. Br). *African Journal of Agronomy ISSN : 2375-1185 Vol. 3 (3)*, pp. 257-258. Available online at www.international scholars journals.org.