

Bioinsecticide effect of *Metarhizium anisopliae* on termite pests *Microtermes lepidus* and *Psammotermes hybostoma*, in the laboratory

Arfang Mafoudji Sonko

Laboratory of Terrestrial Invertebrates Zoology, IFAN Cheikh A. Diop
institute, Cheikh Anta Diop University, Dakar, Senegal
Department of Plant Biology, Faculty of Science and Technology,
Cheikh Anta Diop University of Dakar, Dakar-Fann, Senegal

Dienaba Sall

ISRA Scientific Direction Senegalese Institute for Agricultural Research,
Dakar, Senegal

Mouhamadou Moustapha Ndiaye

Abdoulaye Baïla Ndiaye

Laboratory of Terrestrial Invertebrates Zoology, IFAN Cheikh A. Diop
institute, Cheikh Anta Diop University, Dakar, Senegal

[Doi:10.19044/esj.2025.v21n15p42](https://doi.org/10.19044/esj.2025.v21n15p42)

Submitted: 04 April 2025

Accepted: 14 May 2025

Published: 31 May 2025

Copyright 2025 Author(s)

Under Creative Commons CC-BY 4.0

OPEN ACCESS

Cite As:

Sonko A.M., Sall D., Ndiaye M.M. & Ndiaye A.B. (2025). *Bioinsecticide effect of *Metarhizium anisopliae* on termite pests *Microtermes lepidus* and *Psammotermes hybostoma*, in the laboratory*. European Scientific Journal, ESJ, 21 (15), 42.

<https://doi.org/10.19044/esj.2025.v21n15p42>

Abstract

In Senegal, many studies have shown the economic importance of termite damage to crops. The majority of farmers use dangerous chemicals on human beings, animals, crops, soil, and environment health. Thus, the use of entomopathogenic fungi would be an effective and environmentally eco-friendly alternative to chemical pesticide use. The objectives of this study are to i) carry out infestations in the laboratory, ii) determine the lethal time (LT50) and iii) observe the germination of spores on the corpses of these species. The methodology is based on i) the infestation of 10 workers of *Microtermes lepidus* and *Psammotermes hybostoma* with *Metarhizium* solutions, ii) the monitoring of mortalities, iii) the determination of the lethal time 50 (LT50) and iv) germination of spores on corpses.

For the determination of LT50, the concentrations of 1 g/L and 1.5 g/L gave an LT50 of less than one day in workers of *Microtermes lepidus* and *Psammotermes hybostoma*. For the concentration of 0.5 g/l, the LT50 is 2.5 days for workers of *Psammotermes hybostoma* and less than one day for those of *Microtermes lepidus*. Incubation revealed the appearance of spores on the cadavers, which were given a concentration of 0.5 g/l. This study shows that mortality is related to the concentration of the solution. The higher the concentration, the shorter the LT50 time. The bioinsecticide effect of *Metarhizium anisopliae* is not immediate, resulting in an LT50 of 1 to 2 days, depending on the species.

Keywords: Bioinsecticide, *Metarhizium anisopliae*, termites, laboratory

Introduction

Termites have been considered pests because of the damage they cause to crops and homes (Sib, 2022). Species of the genera *Reticulitermes* and *Coptotermes* (Family Rhinotermitidae) are the major termite pest species in the Americas, Europe and Asia (Pearce, 1997). While Macrotermitidae (especially *Odontotermes*, *Macrotermes* and *Microtermes*) are the predominant termite pest species in Africa and Asia (Pearce, 1997).

In West Africa, and more particularly in Côte d'Ivoire, termites are responsible for damage to rice and maize fields in the sub-Saharan savannah (Akpesse *et al.*, 2008). Similarly, Coulibaly *et al.* (2014) were able to identify the species responsible for the damage to mango tree nurseries in northern Côte d'Ivoire and estimated that termite attacks can cause losses of up to 93.33%.

In Togo, in sugar cane plantations, losses on cuttings caused by termites are estimated at 51% for an economic loss of 29% at harvest (Kotoklo *et al.*, 2011).

In Senegal, studies by Han and Ndiaye (1996) on termite attacks on 10 fruit species revealed attack rates of up to 76.2%. In the regions of Casamance, Saint-Louis, Thiès and Kaolack (Senegal), termites belonging to the trophic groups of lignivores and fungi have been the cause of serious attacks that can lead to the death of certain fruit species (Ndiaye and Han, 2000; 2002; 2006; 2007).

In the department of Tivaouane (Thiès region), the breadbasket of cassava production in Senegal, a study showed the susceptibility of 5 varieties of cassava to attacks by termites belonging to the genera *Macrotermes*, *Odontotermes*, *Amitermes*, *Psammotermes* and *Microtermes* (Faye *et al.*, 2014) and (Faye, 2016).

Several methods of chemical management of these termite pests have been used without success. Thus, in the context of the agroecological

management of these pests, the use of entomopathogenic fungi is an effective and environmentally friendly alternative.

Entomopathogenic fungi have the power to kill or reduce the vigor of the host they infect. *Metarhizium anisopliae* is a mitosporic fungus with asexual reproduction. The classification of *Metarhizium anisopliae* has undergone several revisions in recent years (Benserradj 2014). There are two forms of *Metarhizium* : *Metarhizium flavoviride* and *Metarhizium anisopliae*. The objective of this study is to evaluate the bioinsecticide effect of *Metarhizium anisoplea* in two species of termite pests. Specifically, it's the question of carrying out infestations in the laboratory, determining the lethal time 50 (LT50) and noting the germination of the spores on the corpses. For the purposes of laboratory infestation tests, workers of *Microtermes lepidus* and *Psammotermes hybostoma* are the species selected. Their choice for testing is justified by their frequency in cassava attacks and the extent of the damage that follows.

Methodology

Laboratory environment

The tests were carried out at the Laboratory of Zoology of Terrestrial Invertebrates of the Institut fondamental d'Afrique noire Ch. A. Diop (IFAN-Ch. A. Diop).

In the laboratory, the tests were carried out under conditions of an average temperature of 25.5 °C and an average relative humidity of 56.5%.

Preparation of the fungal solution

The fungal solution was prepared with a strain of *Metarhizium anisopliae* provided by the Directorate of Plant Protection of Senegal. The viability of the strain is 82.33% (Diatta, 2018).

The solution was prepared from 1g of *Metarhizium anisopliae*, 20 ml of distilled water and 2 drops of tween 80. By adding 250 ml of distilled water, the solution is brought to a concentration of 4 g/l.

From this solution, 1 ml was taken and then added to 9 ml of distilled water to evaluate the concentration of conidia in the solution. By depositing 50 microliters of the solution on a Malassez cell, the observation under the LEICA DME microscope at magnification 40 made it possible to evaluate the concentration of fungal spores in the solution, which is $1.19e^6$ spores/ml, using the following formula.

The concentration C of conidia per ml is obtained by the following formula:

$$c = X \times 5.10e^4 c = X \times 5.10e^4$$

X, the average number of conidia counted is obtained by the formula:

$$X = \frac{(a + b + c + d)}{4}$$

a, **b**, **c** and **d** being respectively the number of repetitions performed, corresponding to the number of slides prepared.

Termite infestation tests in the laboratory

Termites

The termites used in the tests were obtained by a bait trap containing cardboard. The traps were buried about 20 cm above the ground.

In the laboratory, petri dishes lined with blotting paper soaked in distilled water were prepared to accommodate the specimens to be tested.

Fungal inocula

Inocula are made up of three concentrations: 0.5 g/l, 1 g/l and 1.5 g/l. The 10 termite workers placed in petri dishes were contaminated by spraying with a micropipette. Each termite received a drop of inoculum. The control boxes contain individuals who have been treated with water. For each treatment, three replicates were performed. The Petri dishes were then placed in a wooden oven.

Test Tracking

Termite monitoring is carried out every day. Dead termites are counted and removed from the boxes and then incubated to control the sporulation of the fungus. The corpses are placed in petri dishes lined with blotting paper moistened with distilled water. After 72 hours of incubation in an oven, the boxes are removed for the observation of termites under the binocular magnifying glass.

Termite mortality rate during testing

The termite mortality rate (TM) corresponding to the ratio is obtained by:

$$TM = \frac{\text{Number of dead individuals}}{\text{Total number of individuals}} \times 100$$

Lethal Time 50 (TL50)

The LT50 corresponds to the time needed to obtain 50% mortality of individuals. After plotting the mortality curve as a function of time, it is obtained by projecting 50% mortality on the x-axis (time).

Statistical analyses

The table of cumulative numbers is used to get an idea of the daily evolution of the number of dead individuals. An ANOVA test with XLSTAT version 2014 is used at the 5% threshold to compare the termite mortality rate between the different treatments used.

Results

Mortality of workers of *Microtermes lepidus*

Statistical tests performed show that there is no significant difference between control termites treated with water and those treated with the 0.5 g/L concentration (Figure 1). Similarly, no difference was observed between termites treated with the 1 g/l and 1.5 g/l concentrations. On the other hand, a significant difference was noted in the evolution of the mortality rate of control termites treated with 0.5 g/l compared to those treated with 1 g/l and 1.5 g/l concentrations.




Figure 1: Mortality during worker testing of *Microtermes lepidus*

Termites treated with 1 g/L and 1.5 g/L did not spore. Sporulation of fungal spores was noted on termites treated with a concentration of 0.5 g/l.

Lethal Time (LT50) for *Microtermes lepidus* Workers

Depending on the concentration, the number of dead termites varies over time. Thus, for termites treated with concentrations of 1 g/l and 1.5 g/l, the LT50 is less than one day. For the 0.5 g/L concentration, the LT50 is 2.5 days (Figure 2).




Figure 2: Evaluation of LT_{50} of different concentrations of *Metarhizium anisopliae* on *Microtermes lepidus* workers in the laboratory

The sporulation of the fungus on the corpses of *M. Lepidus*

The sporulations of fungus, *Metarhizium anisopliae*, are observed on the corpses of termites treated with a concentration of 0.5 g/l. The insects' corpses are colonized by the greenish spore's characteristic of *M. anisopliae* after 11 days of incubation (Figure 3).




Figure 3: Sporulation of *Metarhizium anisopliae* on cadavers

Given the desired effect in the fungus, the transmission of the infestation to the colony by workers, the 0.5 g/L dose, which takes longer to kill termites and which gives evidence of the development of the fungus with sporulation, should be used in termite control tests.

The mortality of workers of *Psammotermes hybostoma*

The control individuals all died after 5 days. Statistically, there were no significant differences between the different concentrations tested (Figure 4).




Figure 4: Mortality during testing of workers of *Psammotermes hybostoma* infested with different concentrations of *Metarhizium anisopliae*

Lethal Time (LT₅₀) for workers of *Psammotermes hybostoma*

For all three concentrations, the LT₅₀ is less than 24 hours. In addition, it is shorter for the 1g/l and 1.5 g/l concentrations of the LT₅₀ than for the 0.5 g/l concentration (Figure 5).




Figure 5: Evaluation of LT₅₀ of different concentrations of *Metarhizium anisopliae*

Sporulation of the fungus on the corpses of *P hybostoma*

Sporulation on corpses is observed after 11 days of incubation only on individuals treated with the concentration 0.5 g/l. *Metarhizium*'s spores cover the entire exoskeleton of the dead insect (Figure 6).



Figure 6: Sporulation of *Metarhizium anisopliae* on a cadaver

Discussion:

The *Metarhizium anisopliae* infestation of termite workers carried out in the laboratory showed that the LT50 varies according to the species. For workers of *Microtermes lepidus* and *Psammotermes hybostoma* exposed to concentrations of 0.5, 1 and 1.5 g/l of LT50, the short lethal time could be explained by several parameters related to the insect, in particular size, but also to the dose received. In this laboratory work, under the same incubation conditions, Diatta (2018) had obtained a 4-day LT50 for the concentrations of 1 and 2 g/l on workers of *Macrotermes subhyalinus* and *Amitermes evuncifer*. The results show that the time taken by *Metarhizium* to kill the insect is a function of the dose applied, as Kpindou *et al.* had pointed out. (2012) but also according to the size of the insect. The smaller the insect, the faster the lethal time and vice versa. The same remarks were made by Kpindou *et al.* (2012) on the different larval stages of *Helicoverpa armigera*, the more advanced the stage, the higher the dose to be applied. Statistical analyses showed that the mortality rates recorded for the 1g/l and 1.5 g/l concentrations are significantly higher than those recorded for the 0.5 g/l concentration and the control. The higher the dose of *Metarhizium anisopliae*, the shorter the lethal time and the higher the mortality rate. These results are similar to those of Diatta (2018), who had higher mortality rates with the high doses and those of Kpindou *et al.* (2012), who also had low levels of pupae and adults of *Helicoverpa armigera* for high concentrations.

During incubation of cadavers, sporulation was observed on individuals tested at the lowest concentration (0.5 g/l) on *Macrotermes subhyalinus* and

Amitermes evancier. These observations were made by Diatta (2018), who recorded sporulations at the lowest concentrations. The strain of *Metarhizium anisopliae* is the same as that used by Diatta (2018). The sporulation of *Metarhizium anisopliae* on cadavers is closely related to the concentration of conidia received by the insect during infection.

Conclusion

The evaluation of the insecticide effect of *Metarhizium anisopliae* in the laboratory has made it possible to understand that mortality is strongly related to the concentration of the solution. For a higher concentration, the mortality rate is higher. The LT_{50} has made it possible to understand that the higher the concentration, the shorter the time. Thus, the bioinsecticide effect of *Metarhizium anisopliae* is not immediate, which gave a LT_{50} of 1 to 2 days depending on the species. The lethal time 50 (LT_{50}) for a concentration of 1 g/l on a target such as *Macrotermes subhyalinus*, which is a large termite, is longer than that of a target such as *Microtermes lepidus* or *Psammotermes hybostoma* which are small species. For the germination of the fungus on the corpse, the lower the concentration of spores, the more the spores have the opportunity to germinate on the corpse. In the future, field trials will be carried out to better assess the real situation.

Acknowledgment

Thanks to Dr. Abdou Aziz Niang, Head of the Laboratory of Terrestrial Invertebrate Zoology of the Institut Fondamental d'Afrique Noire (IFAN Ch. Anta Diop) for his welcome to his laboratory.

Conflict of Interest: The authors reported no conflict of interest.

Data Availability: All data are included in the content of the paper.

Funding Statement: The authors did not obtain any funding for this research.

References:

1. Akpessa A. A., Kouassi P. K., Tano Y., Lepage M. (2008). Impact des termites dans les champs paysans de riz et de maïs en savane subsaharienne (Booro-Borotou, Côte-d'Ivoire). *Sciences & Nature*, 5 (2) ; 121-131.
2. Benserradj O. (2014). Evaluation de *Metarhizium anisopliae* à titre d'agent de lutte biologique contre les larves de moustiques. *Thèse de doctorat 3^{ème} cycle LMD en Biotechnologies, Biologie et Environnement option bioprocédés et biotechnologies, applications mycologiques*, Faculté des Sciences de la nature et de la vie,

- département de Microbiologie, Université Constantine 1, Algérie, 208 p.
3. Coulibaly T, Akpessè A A M, Yapi A, Zirihi G N, et Kouassi K P. (2014). Dégâts des termites dans les pépinières de manguiers du nord de la Côte d'Ivoire (Korhogo) et essai de lutte par utilisation d'extraits aqueux de plantes. *Journal of Animal & Plant Sciences*, 22 (3) : 3455-3468
 4. Diatta C. (2018). Test d'efficacité de deux souches de *Metarhizium anisopliae* (Metchnikoff) Sorokin (*Ascomycotina : Hyphomycetes*) sur des ouvriers de *Macrotermes subhyalinus* (Rambur) (Termitidae, Macrotermitinae) et d'*Amitermes evuncifer* Silvestri (Termitidae, Termitinae) au laboratoire. *Mémoire de Master en phytopharmacie et protection des végétaux ; UCAD, Faculté des sciences et technique, département de Biologie Végétale*, 53 p.
 5. Faye A. (2016). Etude comparative de variétés de manioc (*Manihot esculenta* CRANTZ) selon leur sensibilité aux termites ravageurs des boutures dans le département de Tivaouane (Sénégal) et leurs capacités organogénétiques *in vitro*. *Thèse de Doctorat, spécialité production et protection des végétaux, Faculté des Sciences et Technique, Université Cheikh Anta Diop de Dakar*, 160 p.
 6. Faye A., Kane P. D., Mbaye D. F., Sall Sy D., Sane D. (2014). Study of the cassava varietal sensitivity to termites ravaging cuttings planted in farms in the department of Tivaouane (Senegal). *International Journal of Science and Advanced Technology*, 4 (6):6-16.
 7. Han S. H. & Ndiaye A. B. (1996). Dégâts causés par les termites (Isoptera) sur les arbres fruitiers dans la région de Dakar (Sénégal). *Actes Coll. Insectes Sociaux*, (10) : 111-117.
 8. Kotoklo A. E., Amevoin K., Robert A., Tano Y., Rouland-Lefevre C., Glitho A. I. (2011). Dégâts causés par les termites sur les cannes à sucre au sud du Togo. *Cam. Jour. Biol. Bioch. Sc.*, 19, 1-10.
 9. Kpindou O, K, D, Djegui D A, Glitho I A, Tamò M. (2012). Réponse des stades larvaires de *Helicoverpa armigera* (Hübner) (Lepidoptera : Noctuidae) à l'application de champignons entomopathogènes *Metarhizium anisopliae* et *Beauveria bassiana*. *Biotechnol. Agron. Soc. Environ.* 2012 16(3), 283-293.
 10. Ndiaye A. B. & Han S. H. (2000). L'attaque des arbres fruitiers par les termites dans les vergers de Saint-Louis et de Thiès (Sénégal). *Actes Coll. Insectes Sociaux*, (13) : 127-132.
 11. Ndiaye A. B. & Han S. H. (2002). Attaque des arbres fruitiers par les termites en Casamance (Sénégal) (Isoptera). *Bulletin de la société entomologique de France*, 107 (2) : 193-199.

12. Ndiaye A. B. & Han S. H. (2006). L'attaque des arbres fruitiers par les termites dans la région de Thiès (Sénégal) (Isoptera). *Bulletin de la société entomologique de France*, 111 (1) : 59-64.
13. Ndiaye A. B. & Han S. H. (2007). Les termites (Isoptera) des arbres fruitiers de la région de Kaolack (Sénégal). *Bulletin de l'IFAN Cheikh Anta Diop, Dakar, TLII, série A*, (1-2) : 147-161.
14. Pearce M. J. (1997). Termites Biology and pest management. *CAB International Editeurs*, 172 p.
15. Sib O. (2022). Effet de l'ombrage et de la fiente de poule sur la dynamique des termites et des fourmis dans les cacaoyères de la région de la Nawa (Sud-Ouest, Côte d'Ivoire). Thèse doctorat, Université Jean LOROUGNON GUEDE, 232 p.