

Pregnancy in a Woman With Untreated Vesico-Cloacal Extrophy: A Clinical Case

Roch M'betid-Degana

Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic

Gilles-Davy Kossa-Ko-Ouakoua

Department of Gynecology and Obstetrics, Camp Henri IZAMO Hospital (National Gendarmerie), Bangui, Central African Republic

Jonathan Sabah

Curie Institute, Paris, France

Geniva Gracelia Vanciane M'betid-Degana

Hilary Mattie Kossa Yamenda

Faculty of Health Sciences, University of Bangui, Central African Republic

Antoine Doui-Doumgba

Department of General Surgery, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic

[Doi:10.19044/esj.2026.v22n18p48](https://doi.org/10.19044/esj.2026.v22n18p48)

Submitted: 10 April 2026

Accepted: 26 June 2026

Published: 30 June 2026

Copyright 2026 Author(s)

Under Creative Commons CC-BY 4.0

OPEN ACCESS

Cite As:

M'betid-Degana, R., Kossa-Ko-Ouakoua, G-D., Sabah, J., M'betid-Degana, G.G.V., Kossa Yamenda, H.M., & Doui-Doumgba, A. (2026). *Pregnancy in a Woman With Untreated Vesico-Cloacal Extrophy: A Clinical Case*. European Scientific Journal, ESJ, 22 (18), 48. <https://doi.org/10.19044/esj.2026.v22n18p48>

Abstract

Introduction: Pregnancy in patients with vesicocloacal exstrophy is extremely rare and poorly documented in the literature. **Report case:** We report the case of a 16-year-old adolescent with multiple congenital malformations who was pregnant, referred from a rural health facility without a technical platform and qualified human resources (Batangafo District Hospital) to a facility with limited resources (Sino-Central African Friendship University Hospital Center in Bangui, the capital of the Central African Republic). The patient was successfully surgically managed: a caesarean section was performed, resulting in the cephalic extraction of a live female newborn weighing 2,950 g with an Apgar score of 10/10 at the 1, 5 and 10

minutes of life and the reconstructive surgery for various malformations. The post-operative period was simple and the patient was discharged on day 15.

Conclusion: This rare case of multiple congenital malformations during pregnancy highlights the importance of prenatal follow-up, ultrasound assessment and multidisciplinary management.

Keywords: Multiple congenital malformations; Vesico-cloacal exstrophy; Batangafo; Central African Republic

Introduction

Congenital malformations are a major cause of morbidity and mortality (Sabiri N et al., 2013). Their causes are multiple; they can be due to endogenous anomalies (genetic or chromosomal), but also result from an extrinsic factor affecting embryogenesis: medications (e.g., neuroleptics, antiepileptics), infections, environmental exposure affecting one or both parents (Thomas J et al., 2018). Sometimes, the etiology remains unknown. According to a recent study, child deaths due to congenital malformations are too high in sub-Saharan Africa compared to high-income countries (SciDevNet, 2021), and the Central African Republic (CAR) is no exception. We report the case of a 16-year-old adolescent with multiple congenital malformations, carrying a pregnancy in a context of very limited resources, which we managed in a multidisciplinary manner, and then we will discuss it with data from the literature.

Clinical Case

The patient was a 16-year-old adolescent, unemployed, living in Batangafo in the CAR. She was referred from the Batangafo District Hospital to the Sino-Central African Friendship University Hospital on 28/04/2024 :

- absence of closure of the left abdominal wall (laparochisis) and a congenital umbilical hernia;
- complete absence of the labia majora (total aplasia of the labia majora);
- absence of the vagina except for a small opening providing access to the partial vagina agenesis;
- bladder exstrophy;
- malposition of the urethral meatus in front of the clitoris (epispadias);
- bifid clitoris.

All based on a notion of amenorrhea of 8 months and 2 weeks according to the patient.



Up ↑→ Left

Figure 1. Pregnant polymalformed patient (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)



Figure 2. Ultrasound showing the fetal heart rate (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic)

Family history: The patient is the 4th child in a sibling group of 7, all of whom appear to be in good health. No known hereditary defects have been reported in the family, either on the paternal or maternal side.

Personal history :

- Age at menarche: 13 years;
- Date of last menstrual period unknown;
- Menstrual cycle: irregular (31 days for 5 months then 28 days);
- Primigravida;
- Nulliparous;
- No history of contraceptive use;
- Hospitalized at 7 years old for paraplegia for one week.

The **CLINICAL EXAMINATION** performed on admission on 28/04/2024 finds:

- A good state of consciousness with a Glasgow score of 15/15;
- Good skin and mucous membrane coloration;
- Good hemodynamic condition;
- Weight = 55 kg, Height = 1.62 m => Body Mass Index (BMI) = 20.9 kg/m²;
- An abdomen increased in volume, with a long longitudinal axis, breathing well;
- Active fetal movements well perceived by the mother;
- Uterine height at 33 cm;
- Mobile cephalic presentation;
- A plan of the back on the right;
- An umbilical bulge allowing palpation of certain abdominal organs;
- A pubic anomaly with the absence of the pubic symphysis bones;
- Absence of the labia majora;
- The presence of a protruding suprapubic lesion with visible exposed bladder mucosa with urinary leakage;
- A bifid clitoris;
- Malposition of the urethral meatus in front of the clitoris (epispadias) and partial congenital absence of the vagina with a small vaginal opening providing access to the cervix;
- Fetal heart sounds at 139 beats per minute in the lower right quadrant;
- Vaginal examination is impossible as the vagina is almost nonexistent except for a small opening that barely admits the little finger;
- Rectal examination unremarkable.



Up ← Down →

Figure 3. Patient lying on the operating table with malformations (umbilical hernia, cloacal bladder exstrophy) (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)



Up ↑→ Left

Figure 4: Uro-digestive and genital malformations (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)

The following **paraclinical examinations** were performed:

- Blood group: A / Rhesus: negative;
- Hemoglobin level = 10.6 g/dl;
- C-reactive protein = 4 mg/L;
- HIV 1 and 2 serology: negative;
- Syphilis serology: negative;
- Urine culture (cytobacteriological examination of urine) : Nitrites = positive => urinary tract infection with *Staphylococcus aureus*;
- Obstetric ultrasound: Single intrauterine pregnancy at 36 weeks of amenorrhea, cephalic presentation. Anterior placenta, Grade III. Amniotic fluid normal (Amniotic Fluid Index = 4.3 cm => largest vertical pocket method). Estimated Fetal Weight = 2,700 g;
- Frontal pelvic X-ray: widening of the pubic symphysis bones less than 0.5 cm.

In total: picture of a polymalformative syndrome complicated by a urinary tract infection with *Staphylococcus aureus* associated with anemia in a pregnancy of 36 weeks of amenorrhea. The patient was immediately hospitalized in the department under strict maternal-fetal monitoring with antibiotic treatment and iron supplementation. After three days of treatment, the indication for a prophylactic cesarean section was made because vaginal delivery was practically impossible due to multiple malformations and pelvic diastasis. This cesarean section was performed and allowed the cephalic extraction of a live female newborn weighing 2,950 g with Apgar score of 10/10 at the 1, 5 and 10 minutes of life.



Up ↑ Down →

Figure 5. Painting of the patient (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)



Up ↑ Down →

Figure 6. Extraction of the newborn (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)



Up ↑ Down →

Figure 7. Clamping then cutting of the umbilical cord (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)

The procedure continued with the repair of the laparochisis and the reconstruction of the anterior surface of the bladder. For this, the nearby

abdominal parietal peritoneum was harvested to close it. A capacity of about 30 ml was obtained; an indwelling urinary catheter was then placed, all of which was covered by the aponeurosis. The closure was done with separate stitches.

The procedure continued with the opening of the small opening that constitutes the vagina by enlarging the orifice to create a neo-vagina. A depth of 5 cm, communicating with the urinary catheter in place, was obtained for the evacuation of urine and any potential sexual intercourse.



Up ↑ Down →

Figure 8. Repair of laparochisis and the anterior surface of the bladder (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)



Up ↑ Down →

Figure 9. Skin closure (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)



Up ← Down →

Figure 10. Appearance of the abdominopelvic wall after repair with the presence in yellow of an indwelling catheter (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)



Up ↑ Down →

Figure 11. Appearance of the abdominopelvic wall after repair without the indwelling catheter (Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/04/2024)

The total duration of the procedure was 2 hours and 15 minutes. No anesthetic or surgical incidents were reported. Blood loss was estimated at 400 ml, uncompensated. The patient was transferred to intensive care and then to the ward for post-operative monitoring with the following treatment: injectable Ampicillin 2 g every 8 hours, injectable Metronidazole 500 mg every 8 hours, injectable Gentalline 160 mg every 24 hours, injectable Perfalgan 1 g every 6 hours, and injectable Diclofenac 75 mg every 12 hours for 4 days, then switched to oral. Anti-D immunoglobulin was administered prophylactically. Vital signs were normal; urine output was 1800 ml the next day. Bowel function resumed 24 hours later. Perineal rehabilitation was initiated to allow urinary evacuation. Good wound healing was achieved after 15 days. Therefore, the post-operative course was uneventful.



Up ↑ Down →

Figure 12. Appearance of the abdominal wall after healing (*Photo: Dr. Roch M'BETID-DEGANA and Dr. Gilles-Davy KOSSA-KO-OUAKOUA, Department of Gynecology and Obstetrics, Sino-Central African Friendship University Hospital Center, Bangui, Central African Republic, 30/07/2024*)

The patient was monitored every 15 days for 3 months, then monthly for 3 months. At the end of her six months of follow-up, the patient was discharged. Oral contraception was established after the patient's consent for at least two years, with an absolute contraindication to vaginal delivery during the next pregnancy.

Discussion

1. Age

Our patient was 16 years old. However, there are not enough scientific publications specifically discussing the age of procreation in a woman with multiple congenital malformations, due to the fact that this situation is exceptional and very diverse. Yet, these multiple congenital malformations are often responsible for infertility and complications, both obstetrical and urinary (Amaral PP et al., 2022). We found no cases in the literature of a minor with multiple malformations carrying a pregnancy. Nevertheless, age influences procreation and remains a major factor in fertility regardless of the malformations. The maximum reproductive age is between 20 and 35 years (general concept in Medically Assisted Procreation) (Gaille M, 2019). Fertility is often reduced in a woman with multiple malformations, so the use of Medically Assisted Procreation is often considered in developed countries as an alternative.

2. Mode of delivery

There is almost no scientific article specifically addressing the mode of delivery in a patient with multiple malformations. The mode of delivery is not standardized in a patient with multiple malformations. However, the obstetric literature allows us to establish solid scientific principles based on comparable situations. Thus, our patient underwent a prophylactic cesarean section at eight and a half months, followed by systematic reconstructive surgery. Our approach is justified by the combination of these multiple malformations (pubic diastasis, narrow vaginal opening) preventing the passage of the fetal mass through the birth canal in an immature pelvis (Schmitz T, 2012; Ben-Guigui J et al., 2018).

3. Types of malformations

Our patient presents with:

- The Bladder/Cloacal Exstrophy Complex (BCEC): BCEC encompasses a set of congenital urological anomalies affecting the bladder, urethra, genital organs, and the pelvic musculoskeletal system (Fendereski K and al., 2024). It can impact sexual health and fertility (Bujons A and al., 2016). The diagnosis was delayed in our patient. This situation is not exceptional in Africa (Tshimbayi M and al., 2014). Normally, BCEC is diagnosed during the first trimester of pregnancy. However, in most developing countries, it is detected at birth, as in the case of our patient. This diagnosis still poses significant challenges in rural areas, and management is complicated. This could explain why our patient was not treated for sixteen years. Its association with pregnancy is very rare but not impossible (Surer I and al., 2001).

- Malformations of the genital apparatus, such as urethral meatal atresia, bifid clitoris, congenital atresia of the labia majora, and partial congenital agenesis of the vagina associated with pelvic diastasis: Spontaneous pregnancy may be possible even without reconstructive surgery, as in the case of our patient. However, a case of congenital vaginal agenesis followed by pregnancy has been described in the literature after vaginoplasty (Aimen FJ and al., 2016).

4. Prognosis after reconstructive surgery

Our patient was placed on oral contraception after her consent. However, pregnancy after CEVC repair is not rare (Kalansuriya D and al., 2025). Pregnancy in patients operated on for cloacal exstrophy can present a high risk with obstetric complications (miscarriage, premature birth, premature placental detachment, ...) and urinary complications (urinary infection, urinary obstruction, hydronephrosis, ureteral stricture, and ureteral stones (Deans R and al., 2012) throughout the pregnancy, which highlights the importance of multidisciplinary management during a subsequent pregnancy. Furthermore, the creation of a neovagina for this patient during the procedure will allow not only the restoration of fertility but also the resumption of a "normal" sexual life.

Conclusion

This rare case of vesico-cloacal exstrophy diagnosed late during pregnancy highlights the importance of early diagnosis and multidisciplinary management.

Conflict of Interest: The authors reported no conflict of interest.

Data Availability: All data are included in the content of the paper.

Funding Statement: The authors did not obtain any funding for this research.

Declaration for Human Participants: This study was approved by Dr. Roch M'BETID-DEGAN, gynecologist-obstetrician, Head of the Gynecology-Obstetrics Department of the University Hospital of Sino-Central African Friendship, and the principles of the Declaration of Helsinki were respected.

References:

1. Aimen FJ, Atef Y, Majed G, Radhouane A, Manel M, Monia M et al. Grossesse spontanée après vaginoplastie chez une patiente présentant une aplasie vaginale congénitale. *Journal Asie-Pacifique de la reproduction*. 2016 ;5(4) :351-353).

2. Amaral PP, Ambrosio P, Condeco R, Coelho A, Bello A and Djokovic D. Congenital malformations of the female genital tract: a review of available classification systems. *The International Society for Gynecologic Endoscopy*. 2022;3(3):44-56.
3. Ben-Guigui J, Gauchotte E, Poujois J, Chabot AC, Morel O, Callec R. *La Revue Sage-Femme*. 2018;17(5):208-214.
4. Bujons A, Lopategui DM, Rodriguez N, Centeno C, Caffaratti J, Villavicencio H. Qualité de vie des patientes atteintes du complexe exstrophie vésicale-épispasdias : suivi à long terme. *J Pédiatr Urol*. 2016;12(4):210.e1-6.
5. Deans R., Banks F, Liao L.M., Wood D, Woodhouse C, Creighton SM. Issues reproductives chez les femmes atteintes d'exstrophie vésicale classique : une étude observationnelle transversale. *Am. J. Obstet. Gynecol*. 2012;206.
6. Fendereski K, Schaeffer AJ. Complexe exstrophie vésicale-épispasdias-exstrophie cloacale: caractéristiques, étiologies et données épidémiologiques. *Afr Urol*. Juin 2024;4(se1):S20-S25.
7. Gaille M. L'âge comme critère d'accès à l'assistance médicale à la procréation médicalement assistée. *OpenEditionsJournals*. 2019;50(1):77-96.
8. Kalansuriya D, Roberts J, Britton S, Kimble RM. Grossesse après réparation d'une exstrophie vésicale : un cas clinique. *Représentante de cas Santé des femmes*. 2025 ;18 :47 :e00734.
9. Sabiri N, Kabiri M, Razine R, Kharbach A, Berrada R, Barkat A. Facteurs de risque des malformations congénitales: étude prospective à la maternité Souissi de Rabat au Maroc. 2013;26(4):198-203.
10. Schmitz T. Situations cliniques particulières, maternelles ou foetales, influencant le choix du mode d'accouchement en cas d'antécédent de césarienne. *Journal de Gynécologie Obstétrique et Biologie de la Reproduction*. 2012;41(8):772-781.
11. SciDevNet. Le fardeau des décès d'enfants dus à des anomalies congénitales en Afrique. 2021.
12. Surer I., et al. Pregnancy in woman with bladder exstrophy: a report of two cases and review of the literature. *Journal of Pediatric Surgery*. 2001.
13. Thomas J, Francis F, Fleur D. Malformations congénitales : quels risques en lien avec l'environnement professionnel et extra-professionnel? *ScienceDirect*. 2018;79(3):299.
14. Tshimbayi M, Ndua D, Kazadi C, Shamashanga Kwete L, Bugeme M, Kiopine Mubinda P et al. Exstrophie vésicale: à propos d'un cas diagnostiqué tardivement. *Pan Afr Med J* mars 2014; 17:172.