

CYTOKINES CHANGES ASSOCIATED WITH MENSTRUAL CYCLE IN HIV INFECTED FEMALES AT NAUTH, NNEWI, SOUTH-EAST NIGERIA

Nkiruka .R. Ukibe, PhD

Department of Human Biochemistry, College of Health Sciences, Nnamdi Azikiwe University, Nnewi Campus, Nnewi, Anambra State, Nigeria

Solomon .N. Ukibe, MBBS, MSc

Department of Prosthetics and Orthotics, School of Health Technology, Federal University of Technology Owerri, Nigeria

Obiagel F. Emelumadu, MBBS, FMCPH

Chigozie. O. Ifeadike, MBBS, FMCPH

Department of community medicine, College of Health Sciences, Nnamdi Azikiwe University, Nnewi Campus, Nnewi, Anambra State, Nigeria

Charles.C. Onyenekwe, PhD, FMLSCN

Department of Medical Laboratory Science, College of Health Sciences, Nnamdi Azikiwe University, Nnewi Campus, Nnewi, Anambra State, Nigeria

Joseph .E. Ahaneku, PhD, FSTA(jp), FSeh

Department of Chemical pathology, College of Health Sciences, Nnamdi Azikiwe University, Nnewi Campus, Nnewi, Anambra State, Nigeria

Linus .A. Ilika, Bm. Bch. FmcpH

Department of community medicine, College of Health Sciences, Nnamdi Azikiwe University, Nnewi Campus, Nnewi, Anambra State, Nigeria

Abstract

Background: HIV infection is characterized by hormonal and immunological changes which may grossly affect the reproductive cycle in affected women. **Aim:** To evaluate Cytokine changes in HIV infected women during menstrual cycle. **Materials and methods:** A total of 90 women aged between 15 and 45 years were randomly recruited for the study. 30 of the women were normal healthy seronegative for HIV and served as control. Blood samples were collected under sterile conditions during the follicular and luteal phases of menstrual cycle after due informed consent had been obtained and the samples were analyzed for Cytokines (IL-8, IL-6,

IL-4, and TNF α) using Enzyme Linked Immunosorbent Assay (ELISA) method. **Results:** The Cytokines (IL-8, IL-6, IL-4 and TNF α) were significantly higher at both phases of menstrual cycle in HIV infected women when compared with the Control (P<0.05). **Interpretation and Conclusion:** The study showed significant cytokine changes with some degree of inflammatory reactions in HIV infected women. The implication of these changes within reproductive life of the women is discussed.

Keywords: HIV, Women, Cytokine changes, South East, Nigeria

INTRODUCTION

The Human Immune deficiency Virus/Acquired Immune Deficiency Syndrome (HIV/AIDS) pandemic has become a major public health problem worldwide especially in sub-Saharan Africa where more than 80 percent of all people living with HIV/AIDS reside (UNAIDS 2004; 2012). Reports have shown considerable evidence that the rate at which HIV infection progresses in women is different from that in men (Farzadegan *et al.*, 1998, UNAIDS 2012; USAID, 2012). In Nigeria, about 3.1 million people are living with HIV/AIDS while 58% (1.72 million) are females mostly within reproductive age (UNAIDS 2011; UNAIDS 2012). In Nigeria HIV is a leading cause of morbidity and mortality among women of reproductive age (UNAIDS 2011; NACA 2012).

HIV infection is known to depress the immune system because of the tropic attraction of the virus to the immune cells. Once the immune cells are overwhelmed, the infected host becomes immunocompromised and is then prone to lots of opportunistic infections.

HIV has been known to impact negatively on women's reproductive health (Ikechebelu *et al.*, 2002; Fallahian and Ilkhani, 2006). The negative reproductive impact ranges from menstrual disorders and failure of reproductive function.

Cytokines are low molecular weight extracellular signaling proteins secreted by immune inflammatory cell populations (Desair, 2007). They have been closely associated with ovarian function in females and are believed to be produced locally in the ovulatory follicle where they assist granulosa cell growth and it inhibits their differentiation. They are also believed to stimulate the secretion of ovulation associated substances such as prostaglandins which aid in the ovulatory process (Brannstrom *et al.*, 1995). Cytokines are believed to play a role in menstruation and implantation since they contribute to the defense of the endometrial mucosal epithelium.

Since HIV infection affects all body system including the reproductive system, cytokine changes occur and this may have some undesirable effects on the female reproductive potential.

The present study was therefore designed to evaluate the cytokine changes which are associated with the menstrual cycle in HIV infected females of reproductive age group in Nnewi, South- East, Nigeria.

MATERIALS AND METHODS

The sample population consisted of ninety premenopausal female participants within the age range (15-45years). Thirty participants were apparently healthy controls recruited amongst the Hospital staff. Thirty participants were HIV seropositive females who have not been placed on antiretroviral therapy while the remaining thirty were HIV seropositive female participants who have been placed on antiretroviral therapy for not less than six months. They were recruited at the HIV clinic of Nnamdi Azikiwe University Teaching Hospital Nnewi, Anambra State Nigeria.

After obtaining informed consent of the participants, a well detailed questionnaire was administered to each participant to ascertain their medical and reproductive history. Eight milliliters of venous blood was collected aseptically between 9.00am and 12noon from each participant at follicular (7-13th day) and luteal (21-23rd day) phases of menstrual cycle. All the participants were double screened for malaria parasite and HIV infection using rapid antigen diagnostic techniques for *malaria falciparum* and immunoassay and immunochromatographic methods for HIV screening respectively. The serum content was separated immediately after clot retraction, labeled and stored at -20°C for determination of IL-8, IL-6, IL-4 and TNF α using ELISA (Enzyme Linked Immunosorbent Assay) method.

The ethics committee of NAUTH Nnewi approved the study design and only those who gave their consent were recruited for the study.

Inclusion and Exclusion criteria: only the participant adjudged as HIV stage 2 were recruited for the study. HIV stage 1, stage 3 and stage 4 were excluded, participant with malaria parasite infection as at the time of study were excluded, participants on contraceptives were excluded, women with previous history of infertility prior to the study and participants who were co-infected with tuberculosis were also excluded from the study. Hence the female participants used were those with no prior fertility problems until the existence of HIV infection.

Methods

Antibodies to HIV-1 and HIV-2 in Human Plasma were detected using Abbott Deterimine system, Immunoassay method [(Trinity Biotech UniGold Assay Kit (Trinity Biotech PLC, Ireland)] and immunochromatographic method [(HIV 1 and 2 STAT-PAK Assay kit (Chembio diagnostic system, INC New York, USA)] respectively.

Determination of TNF- α , Interleukin-4, 6 and 8 were done using Enzyme Linked Immunosorbent Assay (ELISA) kits (Glory Science Laboratory USA)

Statistical analysis

The version 16 of SPSS package was used in statistical analysis. The variables were expressed as mean (\pm SD). The student t-test and analysis of variance (ANOVA) and post-hoc (LSD) were used to assess significant mean differences. Graph Pad Prism version 5.03 was used for graph presentations.

RESULTS

Levels of Cytokines (IL-8) at Follicular and Luteal Phases of Menstrual cycle

The mean (\pm SD) plasma IL-8 concentration (pg/ml) dropped significantly at luteal phase (742.5 \pm 197.7, 377.1 \pm 174.2) compared with follicular phase (876.4 \pm 387.2, 550.9 \pm 183.6) of menstrual cycle in HIV seropositive females and HIV seropositive females on ART (P<0.05 respectively). When the mean IL-8 value (pg/ml) at follicular and luteal phases of menstrual were compared between the Control group and Test groups, the mean IL-8 was significantly higher in HIV seropositive females (876.4 \pm 387.2, 742.5 \pm 197.7) and HIV seropositive females on ART (550.9 \pm 183.6, 377.1 \pm 174.2) compared with follicular and luteal values in the Control female subjects (280.1 \pm 47.7, 276.9 \pm 56.3) (P<0.05 respectively).

The post hoc analysis showed significant drop in the mean IL-8 value (pg/ml) at follicular and luteal phases of menstrual cycle in HIV seropositive females on ART (550.9 \pm 183.6, 377.1 \pm 174.2) compared with follicular and luteal values in the HIV seropositive females (876.4 \pm 387.2, 742.5 \pm 197.7) (P<0.05 respectively) (See fig 1).

Levels of Cytokines (IL-6) at Follicular and Luteal Phases of Menstrual cycle

The mean (\pm SD) plasma IL-6 concentration (pg/ml) dropped significantly at follicular phase (474.1 \pm 153.2) of menstrual cycle compared with the luteal phase (584.3 \pm 271.3) in HIV seropositive female subjects (P<0.05). On the other hand, there was no significant difference in the mean IL-6 value (pg/ml) between follicular (224.9 \pm 54.6) and luteal (296.6 \pm 143.7) phases of menstrual cycle in HIV seropositive females on ART (P>0.05). Similarly, there was no significant difference in the mean IL-6 value between follicular (217.6 \pm 64.9) and luteal (204.6 \pm 36.7) phases of menstrual cycle in Control female subjects (P>0.05).

When the mean IL-6 concentration (pg/ml) at follicular and luteal phases of menstrual cycle were compared between the Control group and

Test groups, the mean IL-6 was higher in HIV seropositive females (474.1±153.2, 584.3±271.3) compared with follicular and luteal values in the Control female subjects (217.6±64.9, 204.6±36.7) (P<0.05 respectively).

The post hoc analysis showed significant drop in the mean IL-6 value (pg/ml) at follicular and luteal phases of menstrual cycle in HIV seropositive females on ART (224.9±54.6, 296.6±143.7) compared with follicular and luteal values in the HIV seropositive females (474.1±153.2, 584.3±271.3) (P<0.05 respectively) (See fig 2)

Levels of Cytokines (IL-4) at Follicular and Luteal Phases of Menstrual cycle

The mean (±SD) plasma IL-4 concentration (pg/ml) in HIV seropositive females significantly dropped at luteal (354.9±207.7) compared with follicular (497.6±216.1) phase of menstrual cycle (P<0.05). But the mean IL-4 value (pg/ml) dropped significantly at follicular (527.2±231.3) compared with luteal phases (660.2±254.2) in HIV seropositive females on ART (P<0.05). The mean IL-4 value (pg/ml) dropped significantly at follicular phase (210.7±71.2) compared with luteal phase (334.8±76.5) in Control female subjects (P<0.05).

When the mean IL-4 concentration (pg/ml) at follicular and luteal phases were compared between the Control group and Test groups, the mean IL-4 was significantly higher in HIV seropositive females (354.9±207.7, 497.6±216.1) and HIV seropositive females on ART (527.2±231.3, 660.2±254.2) compared with follicular and luteal values in the Control female subjects (210.7±71.2, 334.8±76.5) (P<0.05 in each case).

The post hoc analysis showed significantly higher mean IL-4 value (pg/ml) at follicular and luteal phases of menstrual cycle in HIV seropositive females on ART (527.2±231.3, 660.2±254.2) compared with follicular value in HIV seropositive females (354.9±207.7, 497.6±216.1) (P<0.05 respectively) (See fig 3).

Levels of Cytokines (TNFα) at Follicular and Luteal Phases of Menstrual cycle

The mean (±SD) plasma TNFα concentration (pg/ml) dropped significantly at luteal phase (788.2±191.7) compared with follicular phase (949.6±335.7) of menstrual cycle in HIV seropositive female subjects (P<0.05). There was no significant difference in the mean TNFα value (pg/ml) between follicular (483.6±160.0) and luteal (519.2±177.8) phases of menstrual cycle in HIV seropositive females on ART (P>0.05). The mean TNFα value (pg/ml) dropped significantly at follicular phase (211.8±57.6) compared to luteal phase (333.0±72.2) of menstrual cycle in Control female subjects (P<0.05).

When the mean TNF α concentration (pg/ml) at follicular and luteal phases of menstrual cycle were compared between the Control group and Test groups, the mean TNF α was significantly higher in HIV seropositive females (949.6 \pm 335.7, 788.2 \pm 191.7) and HIV seropositive females on ART (483.8 \pm 160.0, 519.2 \pm 177.8) compared with follicular and luteal values in the Control female subjects (211.8 \pm 57.6, 333.0 \pm 72.2) (P<0.05 in each case).

The post hoc analysis showed significant drop in the mean TNF α concentration (pg/ml) at follicular and luteal phases of menstrual cycle in HIV seropositive females on ART (483.8 \pm 160.0, 519.2 \pm 177.8) compared with follicular and luteal values in HIV seropositive female subjects (949.6 \pm 335.7, 788.2 \pm 191.7) (P<0.05 respectively) (See Fig 4)




Fig 1: Comparison of mean (\pm SD) plasma levels of IL-8 in Test groups and Control group at Follicular and luteal phases of menstrual cycle




Fig 2: Comparison of mean (±SD) plasma levels of IL-6 in Test groups and Control group at Follicular and luteal phases of menstrual cycle




Fig 3: Comparison of mean (±SD) plasma levels of IL4 in Test groups and Control group at Follicular and luteal phases of menstrual cycle




Fig 4: Comparison of mean (\pm SD) plasma levels of TNF α in Test groups and Control group at Follicular and luteal phases of menstrual cycle

DISCUSSION

The study showed significantly higher levels of IL-8 and TNF α in HIV seropositive females and HIV seropositive females on ART at follicular phase compared to luteal phase of menstrual cycle. Cytokines including TNF α and IL-8 are secreted by activated monocytes. It is worthy to note that monocytes populations are reduced in diseased states such as HIV infection indicating reduced cellular immunity which is one of the problems in these subjects. The increased secretion of these cytokines therefore may be due to hyperstimulation of the surviving monocytic cells. It has been reported that endotoxin stimulated monocytes of women in the luteal phase produced more TNF α when compared to follicular phase (Brannstrom *et al.*, 1999; Bouman *et al.*, 2001). IL-8 was also found to be significantly elevated at the follicular phase compared with the luteal phase (Al- Harthi, 2001) which implied increased inflammatory response and enhanced cellular immunity in this phase and this may have been induced by reduced progesterone and estrogen production. This report is consistent with the observation in the present study.

The significantly higher level of TNF α in HIV seropositive female subjects at the follicular phase when compared with the luteal phase is inconsistent with findings in healthy women. It has been documented that in apparently healthy women, TNF α is generally higher at the luteal phase during which the population of activated monocytes, which secrete TNF α , are reported to be higher (Marthur *et al.*, 1979; Northern *et al.*, 1994; Brannstrom *et al.*, 1999). This is probably due to the fact that the luteal phase is the time when fertilization takes place and the fertilized ovum requires protection from infective agents hence the increased pro-inflammatory activity (Norman, 2001).

The significantly higher level of IL-6 at the luteal phase compared with the follicular phase in the present study is comparable with the previous reports done in developed countries (Konecna *et al.*, 2000). IL-6 is said to stimulate B-lymphocytes (thereby enhancing humoral immunity) and T-lymphocytes differentiation and activate macrophages and NK-cells (thereby enhancing cellular immunity). It is also said to possess anti-inflammatory properties (Marijke *et al.*, 2011). Since IL-6 production is said to be decreased by estrogen, the low estrogen production associated with hypogonadism in HIV subjects result in increased IL-6 production since sex hormones perform immunoregulatory functions.

Reports of IL-6 levels during the menstrual cycle phases have been controversial. Whereas Angstwurm *et al.*, (1997) reported higher levels of IL-6 during the follicular phase of menstrual cycle, Abrahamsen *et al.*, (2003) reported no difference while Schwarz *et al.*, (2000) reported decreased IL-6 in the follicular phase during the menstrual cycle. It has been reported that during the luteal phase of menstrual cycle, the immune response is shifted toward a Th2-type response and this was found to correspond to increased levels of progesterone and 17 β -E₂ (Marijke *et al.*, 2000). This account for the anti-inflammatory immune response associated with IL-6 production at this phase and reduced cellular immunity.

The significantly high levels of IL-8, IL-6, IL-4 and TNF α in HIV seropositive females and HIV seropositive females on ART compared to Control females at both follicular and luteal phases of menstrual cycle also showed higher degree of inflammation in diseased subjects when compared to healthy people. Excessive production of some cytokines for instance TNF α has been associated with problems such as fever or even tumor formation hence certain neoplastic conditions such as Kaposi sarcomas are said to be common in chronic HIV patients (Locksley *et al.*, 2001). The increased IL-4 may be due to anti-inflammatory effects of estrogen thereby promoting Th2 immune response. Moreover, HIV is a chronic inflammatory disease and is said to induce a Th1-type immune responses which are characterized by the production of pro-inflammatory cytokines some of

which have been associated with harmful effects such as fever and tumors formation (Chowdbury *et al.*, 2010). The implications of elevated cytokines on endocrine and metabolic functions have been previously reported (Hashimoto *et al.*, 1994; Gartner, 2009) and this has far reaching effects on menstrual and reproductive functions of the affected women (O' Brien *et al.*, 2007).

However, the significantly reduced levels of IL-8, IL-6, IL-4 and TNF α in HIV seropositive females on ART compared to their counterparts without treatment showed significant reduction of inflammation and perhaps some level of improvement in the reproductive function in these subjects due to a reduction in viral load (AL-Harathi *et al.*, 2001). This signifies beneficial effects of treatment which results in significant restoration of the cellular immunity and reduction of inflammation in these patients. This has been previously reported (Sachdeva *et al.*, 2010). The fluctuations in the levels of cytokines have been reported to be related to hypogonadal function as has been discussed previously (Marijke *et al.*, 2000).

The significantly increased levels of the anti inflammatory cytokine (IL-4) in HIV seropositive females on ART especially at the luteal phase were consistent with the value observed in Control females. These signify a reduction or modulation of inflammatory activities and humoral immunity which is a Th2 type of response. IL-4 is also secreted by activated immune cells such as monocytes. It has been reported that variations in progesterone and estrogen (sex hormones) induced by HIV infections drive the immune response to either a Th1 type or Th2 type response depending on the concentration of these hormones. This leads to the exaggeration of either pro-inflammatory cytokines such as TNF α or anti-inflammatory cytokines such as IL-4 (Marijke *et al.*, 2000). It has also been reported that cytokines are elevated in systemic diseases and this can directly inhibit the ovary in females (Marijke *et al.*, 2000).

The present study therefore, concludes that there were cytokine changes with some degrees of inflammatory reactions in HIV infected females.

References:

- Abrahamsen B, Stilgren LS, Rettmer E, Bonnevie-Nielsen H. Effects of natural and artificial menstrual cycle on the production of osteoprotegerin and the bone resorptive cytokines IL-1 β and IL-6. *Calcif Tissue International* **2003**; 72(1):18-23.
- Al-Harathi L, Wright DJ, Anderson D, Cohen M, matityalu D, Lohn J, CU-llvin S, Burns Reichel derfer P, Levis S, beekner S, Kovacs A, Landay A. The impact of the ovulatory cycle on cytokine production: Evaluation of

systemic, cervicovaginal, and salivary compartments. *Journal of Interferon and Cytokine Research* **2004**; 20(8): 719-724.

Al-Harathi L. A menstrual cycle pattern for cytokine levels exists in HIV-positive women: implication for HIV vaginal and plasma shedding. *Acquired Immune Deficiency Syndrome* **2001**; 15(12): 1535-1543.

Angstwurm M, Gartner R, Zeigler-Heitbroek H. Cyclic plasma IL-6 levels during normal menstrual cycle. *Cytokine* **1997**; 9(4): 370-347.

Bouman A, Moes H, Heineman MJ, De Leji LF, Faas MM. Cytokines production by natural killer lymphocytes in follicular and luteal phase of the ovarian cycle in humans. *American Journal of Reproductive Immunology* **2001**; 45(1):130-134

Brannstrom M, Fridan BE, Jasper M, Norman RJ. Variations in peripheral blood levels of immunoreactive tumour necrosis factor- α (TNF- α) throughout the menstrual cycle and secretion of TNF α from the human corpus luteum. *European Journal of Obstetrics and Gynaecology and Reproductive Biology* **1999**; 83(2): 213-217.

Chirgwin KD, Feldman F, Muneyyirci-Delale O. Menstrual function in Immunodeficiency Virus-infected women without Acquired Immunodeficiency Syndrome. *Journal of Acquired Immune Deficiency Syndrome and Human Retrovirology* **1996**; 12(4): 489-494.

Chowdhury RG, Pain SK, Bhattacharjee B, Chatterjee S. Infestation of Endometrium by *Mycobacterium Tuberculosis Bacilli*-cause of Reproductive Failure. *Al Meen Journal of Medical Sciences* **2010**; 3(4): 322-331

Desai P. (2007). Cytokines in Obstetrics and Gynecology. *Journal of Obstetrics and Gynecology* 237(3):197-200.

Ellerbrock TV, Wright TC, Bush TJ, Dole P, Brudney K, Chiasson MA. Characteristics of menstruation in women infected with Human Immunodeficiency Virus. *Obstetrics and Gynecology* **1996**; 87(6): 1030-1034.

Fallahian M, Ilkhani M. Menstrual Disorders in Nongenital Tuberculosis. *Infectious Disease in Obstetrics and Gynecology* **2006**; 18452:1–3.

Farzadegan H, Hoover Dr, Astemborski J. Sex differences in HIV-1 viral load and progression to AIDS. *Lancet* **1998**; 352(12): 1510-1514.

Gartner R. Selenium and thyroid hormone axis in critical ill states: an overview of conflicting view points. *Journal of Trace Elements and Medical Biology* **2009**; 23(2):71-74

Grinspoon S, Corcoran C, Kareem M, Biller BMK, Askari H, Wang E, Hubbard J, Anderson EJ, Basgoz N, Heller HM, and Klibansky A. Body Composition and Endocrine function in Women with Acquired Immunodeficiency Syndrome Wasting. *Journal of Clinical Endocrinology and Metabolism* **1997**; 82(5):1332-1336.

- Harlow SD Schuman P, Cohen M, Ohmit SE Cu-Uvin S, Lin X, Anastos K, Burns D, green blath R, Minkoff A, Muderspach L, Rompalo A, Warren D, Young MA Klein RS. Effect of HIV infection on menstrual cycle length. *Journal of Acquired immune Deficiency Syndrome* **2000**; 24(1):68-75
- Hashimoto H, Igarashi N, Yachie A, Miyawaki T, Sato T. The relationship between serum levels of interleukin-6 and thyroid hormone in children with acute respiratory infection. *Journal of Clinical Endocrinology and Metabolism* **1994**; 78(2):288-291
- Hutchinson J, Murphy M, Harries R, Skinner CJ. Galactorrhoea and hyperprolactinaemia associated with protease inhibitors. *Lancet* **2000**; 356: 1003-1004.
- Ikechebelu JI, Ikegwuonu SC, Joe-Ikechebelu NN. HIV Infection and Sexual behaviours among infertile women in Southwestern, Nigeria. *Journal of Obstetrics and Gynaecology* **2002**; 22(3): 306-307.
- Konecna L, Yan MS, Miller LE, Scholmerich J, Falk W, Straub RH. Modulation of IL-6 production during the menstrual cycle in vivo and in vitro. *Brain Behaviour and Immunology* **2000**; 14(1): 49-61
- Locksley RM, Killeen N, Lenardo MJ. "The TNF and TNF receptor superfamilies: integrating mammalian biology". *Cell* **2001**; 104 (4): 487–501.
- Marijke F, Paul de Vos, Barbro M. Sex Hormones and Immunoregulation. Available at:
<http://brainimmune.tumblr.com/post/68872756667/sex-hormones-and-immunoregulation-marijke-faas> **2011**.
- Marijke F, Annechien B, Henk M, Maas JH, Loe L, Gerard S. The immune response during the luteal phase of the ovarian cycle: a Th2-type response? *Fertility and Sterility* **2000**; 74(5):1008-1013.
- Mathur S, Mathur RS, Goust JM, Williamson HO, fudenberg HH. Cyclic variations in white cell subpopulations in the human menstrual cycle: correlations with progesterone and estradiol. *Clinical Immunology and Immunopathology* **1979**; 13(3):346-353
- Merrill JE, Koyanagi Y, Chen ISY. Interleukin -1 and tumor necrosis factor α can be induced from mononuclear phagocytes by human immunodeficiency virus type1 binding to the CD4 receptor. *Journal of Virology* **1989**; 63(1):4404-4408
- National Agency for the Control of AIDS (NACA). Federal Republic of Nigeria, Global AIDS Response Country Progress Report Nigeria. Country Progress Report **2012**.
- Norman J. *Jeffcoate's Principles of Gynaecology*. 5th ed. Ansari Nagar, New DELHI, India. **2001**; 60-119.
- Northern AL, Rutter SM, Peterson CM. Cyclic changes in the concentrations of peripheral blood immune cells during the normal menstrual cycle.

Proceedings of Society of Experimental Biology and Medicine **1994**; 207(3):81-88

O' Brien SM, Fitzgerald P, Scully P, Landers A, Lucinda V, Scott Dinan TG. Impact of Gender and Menstrual cycle phase on Plasma cytokines concentrations *Neuroimmunomodulation* **2007**; 14(1): 84-90.

Panda S, Singh AS, Jha Vandana. The concepts and Consequences of Early Ovarian Ageing: A Caveat to Women's Health. *J. Reprod Infertil* **2013**; 14(1):3-7

Sachdeva RK, Wanchu A, Abagga R, Malla N, Snaima M. Effects of non nucleoside reverse transcriptase inhibitors on cytokine, chemokine and immunoglobulin profiles in serum and genital secretions of HIV-infected women. *Journal of Interferon and cytokine Resource* **2010**; 30(5): 299-310

Schwartz LT, ST Louis Y, Wu R, Wiznia A, Rubinstein A, Saenger P. Endocrine function in children with HIV infection. *American Journal of Disease and child* **1991**; 145(3): 330-333.

Shah PN, Smith JR, Wells C, Barton SE, Kitchen VS, Steer PT. Menstrual Symptoms in women infected by the Human Immunodeficiency virus. *Obstetrics Gynecology* **1994**; 83(4): 397-400

Tracey KJ, Cerami A. Metabolic responses to cachectin/TNF: A Brief review. *Annals of Academic of Science* **1990**; 587 (3):325-331

UNAIDS 'World AIDS Day Report' Nov 2011

UNAIDS 'Global Report: Annexes' **2012**

UNAIDS 'Women Out Loud: How Women Living with HIV Will Help the World End AIDS' **2012**

UNAIDS/UNFPA/UNIFEM 'Women and HIV/AIDS: Confronting the crisis' **2004**.

USAID/Nigeria. HIV/AIDS and Tuberculosis **2012**.